

LIBRARY
OF THE
UNIVERSITY
OF ILLINOIS

590.5

FI

v. 35

cop. 3

NATURAL HISTORY
SURVEY

U.S.
N.M.
3

CHECK LIST
OF
NORTH AMERICAN WATER-MITES

RODGER D. MITCHELL

FIELDIANA: ZOOLOGY
VOLUME 35, NUMBER 3
Published by
CHICAGO NATURAL HISTORY MUSEUM
AUGUST 31, 1954

CHECK LIST
OF
NORTH AMERICAN WATER-MITES

RODGER D. MITCHELL

Department of Zoology, University of Vermont

FIELDIANA: ZOOLOGY

VOLUME 35, NUMBER 3

Published by

CHICAGO NATURAL HISTORY MUSEUM

AUGUST 31, 1954

THE LIBRARY OF THE

SEP 20 1954

UNIVERSITY OF ILLINOIS

PRINTED IN THE UNITED STATES OF AMERICA
BY CHICAGO NATURAL HISTORY MUSEUM PRESS

90.5
I
35³
p. 3

North American Water-Mites

INTRODUCTION

Fewer than fifty species of water-mites were known from North America in 1900. Now more than two hundred and seventy species are known, largely through the work of Dr. Ruth Marshall. During her forty years' study of the group Miss Marshall amassed a collection containing abundant material of nearly every species known from this continent. In addition to the type material of her own species, the Marshall Collection contains most of the Wolcott types, paratypes from Lavers, and a few types of species described from North America by Lundblad, Koenike, and Karl Viets. This collection was deposited in Chicago Natural History Museum in 1945. A listing of the types in the Marshall Collection was started and, as the work progressed, it was apparent that with a little expansion a full listing of the North American water-mites could be prepared. The species represented by type material in the Marshall Collection are indicated by the designation "[Marshall Collection]" following the citation. A detailed listing of the types is not possible at present for two reasons. Types of the first few species described by Miss Marshall were destroyed in a fire and subsequent neotype selections have not been published. In most of the descriptions, the type series was not clearly indicated and a formal designation of the holotype was not made. Further clarification of the type material should be undertaken through revisionary studies.

The synonymies in the check list represent, as far as is possible to ascertain, the views of Miss Marshall as expressed in her papers, and, in a few cases, in her notes. Subgenera were rarely indicated by her, hence they can be listed in only a few cases. Furthermore, most authors have defined the subgenera in the course of limited faunistic studies, with the result that general usage of subgenera is difficult. Also, the understanding of the Nearctic fauna is so limited that the use of subgenera is not particularly helpful in most cases. The subgenera of *Hydrachna* and *Arrenurus* are obvious exceptions. It is suggested (Lundblad, 1944) that intergradation between the subgenera *Truncaturus*, *Megaluracarus*, and *Micruracarus* of the

genus *Arrenurus* is a basis for suppressing the use of these categories. The definition of the subgenera is, in respect to the few marginal forms, clearly arbitrary. Regardless of the decision in this matter, it is obvious that each of the subgenera indicates a phyletic line and is of great practical value in breaking up a genus of tremendous size.

In a few cases there is uncertainty about the selection of generotypes, a question on which I do not feel qualified to pass judgment. Current usage by European workers has been followed in these cases. Viets (1949) gives a full discussion of the two major name changes affecting the species considered here. *Atractides* must be applied in place of *Megapus* and the species that have been included in *Atractides* must be referred to under the name *Torrenticola*. This unfortunate change calls for changing the family name Atractideidae to Torrenticolidae. The genus *Acercus* must now be referred to under the name *Tiphys*.

Families of water-mites have been defined entirely by European investigators. The system adopted today was outlined in the papers of Thor, but it has been modified and developed through the research of Viets and Lundblad. Lundblad's classification (1941b) has been followed here because it is the most recent complete listing of the genera. It differs only slightly from that of Karl Viets (1936).

The forty families of water-mites, although closely related, cannot be assigned to an inclusive taxonomic category. Many authors have chosen to cite Hydrachnellae or Hydracarina (Latinized names for water-mites) in the same fashion as a taxonomic category. Since there is general agreement that the group is an ecological and not a phyletic one, such a practice seems illogical and misleading. The various systems for grouping the water-mites with the prostigmatid mites suffer from the lack of any critical information on the internal and skeletal anatomy.

I am aware that much critical study is necessary to clarify the status of many of the groups listed below. Furthermore, many genera known to occur in North America are not yet recorded in the literature. Considering these matters, this check list is offered only as a summary of our published knowledge of the North American water-mites.

The preparation of this list has been greatly facilitated by the kind consideration and helpful criticism of Rupert L. Wenzel and Henry S. Dybas. I am indebted to David R. Cook for his help in reviewing and checking the list.

HISTORY

Before the latter part of the nineteenth century there were only a few publications on our water-mite fauna, all inadequate. The first usable paper, published by Koenike in 1895, gave identifications and descriptions of a surprisingly good collection of water-mites made by J. B. Tyrrell. Around the turn of the century Robert H. Wolcott published a few papers. His key to the genera of the world (1905) was later abridged and included in *Fresh-Water Biology* (1918). The classification suggested in this work is very much out of date due to the subsequent work of various European students.

Between the years 1903 and 1946 Ruth Marshall published forty-one papers. Most of these papers deal with the still-water fauna of the mid-western states. Marshall's descriptions and illustrations form a workable basis for the identification of our water-mites, but it is regrettable that the results of these studies have never been assembled in the form of larger monographs or revisions. Two other significant papers have been published: a revision of the genus *Hydrachna* by Lundblad (1934), and an article by Lavers (1945) on the *Arrenurus* of the state of Washington. Only a few brief descriptive papers on the North American water-mite fauna have appeared since 1900.

Baker and Wharton (1952) have presented a key to, and synopses of, the water-mite families of the world. There are a few inconsistencies in the presentation, because the key, translated from Karl Viets (1936), has not been modified to conform with the classification followed in the family synopses, which were taken from Vitzthum (1940-43).

The most recent publication in this country is that of Pennak (1953), whose key (apparently taken from Wolcott, 1918) is preceded by an account of the biology of water-mites that is remarkably well done, considering the limitations of space. Omissions make the key of little value to the specialist, but probably serve to make the account a comprehensible and reasonably accurate introduction for the non-specialist who deals chiefly with the common genera.

GENERAL WORKS

None of the keys published in this country (Wolcott, 1905 and 1918; Pennak, 1953) are sufficiently complete for advanced studies of water-mites. Because of the uniformity, at least in generic representation, of the Holarctic fauna the excellent monographs on

the European fauna (Viets, 1936; Motas, 1928; and Lundblad, 1927) are indispensable basic references for any advanced work on the North American water-mite fauna. The volumes published by the Ray Society (Soar and Williamson, 1925-29), though lacking in some critical detail, are frequently helpful in supplying general information.

One meets with considerable difficulty in working with the mites that occur in our streams. Most of the genera are little known and many are not Holarctic; therefore, there is no single work that can be used for identifying these forms.

Attention should be called to certain works as sources of special information. The remarkable morphological diversity of water-mites is clearly demonstrated in the work of Motas (1928). Since 1928, Motas and his associates have done a creditable job of describing the subterranean fauna of the Balkans. Most of the work is summarized by Szalay (1949), although Motas' continued researches have added more forms since that summary. In addition to the fine treatments on the European fauna mentioned earlier, there are two other important faunistic studies, Karl Viets' study of the water-mites of Sumatra, Java, and Bali (1935) and the work of Lundblad (1941-44) on the South American fauna, which is a magnificently illustrated monograph. The extensive work on the African fauna has been reviewed and listed by Karl Viets (1953).

Family HYDROVOLZIIDAE Thor

Hydrovolziidae Thor, 1905, Zool. Anz., 28: 509.

Genus *Hydrovolzia* Thor

Generotype: *Polyxo placophora* Monti.

Hydrovolzia Thor, 1905, op. cit., pp. 506-507.

gerhardi Mitchell, 1954, Chicago Acad. Sci. Nat. Hist. Misc., no. 134, pp. 3-6, figs. 1-3. [C.N.H.M. Collection]

placophora Monti, 1905, Rend. Ist. Lomb. Sci. Lett., (2), 38: 170-174, pl. 1, fig. 1, pl. 2, figs. 2-14.

Family HYDRACHNIDAE Leach

Hydrachnides Leach, 1815, Trans. Linn. Soc. London, 11: 399-400.

Genus *Hydrachna* Müller

Generotype: *Hydrachna cruenta* Müller.

Hydrachna Müller, 1776, Zool. Danicae Prodrumus, p. 188.

Subgenus *Hydrachna* s. str.

cruenta Müller, 1776, Zool. Danicae Prodrömus, p. 190.

americana Marshall, 1929, Univ. Toronto Stud., Biol. Ser., no. 33, p. 63, pl. 3, figs. 25-27. [Marshall Collection]

var. *diminuata* Lundblad, 1934, Ark. Zool., 28A, (3), p. 10, figs. 4a-4d.

magniscutata Marshall, 1927, Trans. Amer. Micros. Soc., 46: 271-272, pl. 7, figs. 8-11. [Marshall Collection]

var. *reducta* Lundblad, 1934, Ark. Zool., 28A, (3), pp. 13-14, figs. 6a-6e, pl. 1, fig. 8, pl. 2, fig. 16.

var. *separata* Lundblad, 1934, op. cit., p. 15, figs. 5e-5h, pl. 1, fig. 5, pl. 2, fig. 15.

Subgenus *Scutohydrachna* Viets

Generotype: *Hydrachna dorsoscutata* Viets, 1933, Zool. Anz., 103: 162-164, figs. 1-3.

Scutohydrachna Viets, 1933, Zool. Anz., 103: 162.

Chitohydrachna Habeeb, 1950, Natural. Canad., 77: 116.

crenulata Marshall, 1930, Trans. Wis. Acad. Sci., 25: 247, pl. 5, figs. 6-7. [Marshall Collection]

?ennishonensis Habeeb, 1950, Natural. Canad., 77: 116-117, figs. 7-11. [C.N.H.M. Collection]

hutchinsoni Lundblad, 1934, Ark. Zool., 28A, (3), pp. 20-21, figs. 10a-10e.

rotunda Marshall, 1930, Trans. Wis. Acad. Sci., 25: 246-247, pl. 5, figs. 4-5. [Marshall Collection]

Subgenus *Diplohydrachna* Thor

Generotype: *Acarus globosa* Geer, 1778, Mem. hist. ins., 7: 146-147, pl. 9, figs. 11-12.

Diplohydrachna Thor, 1916, Rev. Ruisse Ent., 16: 46-47.

conjecta Koenike, 1895, Rev. biol. nord. Fr., 7: pp. 145-146, pl. 8, figs. 9-11.

microscutata Marshall, 1929, Univ. Toronto Stud., Biol. Ser., no. 33, p. 65, pl. 3, figs. 28-30. [Marshall Collection]

hungerfordi Lundblad, 1934, Ark. Zool., 28A, (3), pp. 23-25, figs. 13a-13f, 18-19.

Subgenus *Tetrahydrachna* Lundblad

Generotype: *Hydrachna miliaria* Berlese.

Tetrahydrachna Lundblad, 1934, Ark. Zool., 28A, (3), p. 30.

miliaria Berlese, 1888, Bull. Soc. Ent. Ital., 20: 219.

bilunata Marshall, 1931, Trans. Wis. Acad. Sci., 26: 314, pl. 8, figs. 26-27. [Marshall Collection]

Subgenus *Rhabdohydrachna* Viets

Generotype: *Hydrachna comosa* Koenike, 1896, Zool. Anz., 19: 359-360.

Rhabdohydrachna Viets, 1931, op. cit., 93: 184.

- canadensis* Marshall, 1929, (female only), Univ. Toronto Stud., Biol. Ser., no. 33, pp. 63-64, pl. 3, figs. 21, 22, 24. [Marshall Collection]
- geographica* Müller, 1776, Zool. Danicae Prodrum, p. 190 (Palearctic).
 form americana Lundblad, 1934, Ark. Zool., 28A, (3), pp. 30-32, figs. 16a-16e.
- hesperia* Lundblad, 1934, Ark. Zool., 28A, (3), pp. 38-41, figs. 21a-21d, pl. 2, fig. 13.
- marshallae* Lundblad, 1934, op. cit., pp. 34-35, 18a-18f. [Marshall Collection]
 canadensis Marshall, 1929, (male only), Univ. Toronto Stud., Biol. Ser., no. 33, pp. 63-64, pl. 3, figs. 20, 23. [Marshall Collection]
- stipata* Lundblad, 1934, Ark. Zool., 28A, (3), pp. 35-38, figs. 19a-19g, figs. 20a-20c, pl. 2, figs. 9-11.

Family LIMNOCHARIDAE Kramer

Limnocharidae Kramer, 1877, Arch. Naturg., 43, (Bd. 1), pp. 242-244.

Genus *Limnochares* Latreille

Generotype: *Acarus aquaticus* Linnaeus.

Limnochares Latreille, 1796, Précis des caractères des insectes, p. 181.

Subgenus *Limnochares* s. str.

aquatica Linnaeus, 1758, Syst. nat., p. 617.

Subgenus *Cyclothrix* Wolcott

Generotype: *Limnochares crinita* Koenike, 1899, Abh. Senckenb. naturf. Ges., 21: 313-318, pl. 21, figs. 23-29.

Cyclothrix Wolcott, 1905, Trans. Amer. Micros. Soc., 26: 185.

americana Lundblad, 1941, Zool. Anz., 133: 155.

natans Lavers, 1941, Univ. Wash. Pub., Biology, 12: 3-6, pl. 1, figs. 1-3.

?acadiensis Habeeb, 1950, Natural. Canad., 77: 114, 116, figs. 1-3.
 [C.N.H.M. Collection]

Family EYLAIDAE Leach

Eylaides Leach, 1815, Trans. Linn. Soc. London, 11: 399.

Genus *Eylais* Latreille

Generotype: *Hydrachna extendens* Müller, 1776, Zool. Danicae Prodrum, p. 190.

Eylais Latreille, 1796, Précis des caractères des insectes, pp. 182-183.

abitibiensis Marshall, 1929, Univ. Toronto Stud., Biol. Ser., no. 33, pp. 60-62, pl. 1, figs. 1-4. [Marshall Collection]

extendens Müller, 1776, Zool. Danicae Prodrum, p. 190.

bakeri Marshall, 1946, Trans. Wis. Acad. Sci., 36: 357, pl. 4, figs. 37-38.
 [Marshall Collection]

desecta Koenike, 1897, Abh. naturw. Ver. Bremen, 14: 288-289.

falcata Koenike, 1897, op. cit., p. 288.

gibberipons Viets, 1910, op. cit., 20: 171-173, figs. 5-6.

hirsutipalpis Marshall, 1946, Trans. Wis. Acad. Sci., 36: 357-358, pl. 4, figs. 42-43. [Marshall Collection]

infundibulifera Koenike, 1897, Abh. naturw. Ver. Bremen, 14: 284-285.

marshallae Koenike, 1912, Trans. Royal Canad. Inst., 9: 286-288, pl. 1, figs. 8-11, pl. 2, fig. 10.

rimosa Piersig, 1899, Zool. Anz., 22: 65-66, fig. 6.

triangulifera Koenike, 1897, Abh. naturw. Ver. Bremen, 14: 289-290.

infundibulata Lundblad, 1941, Zool. Anz., 133: 155.

robusta Marshall, 1946, Trans. Wis. Acad. Sci., 36: 358, pl. 4, figs. 44-46. [Marshall Collection]

Family PROTZIIDAE Viets

Protziidae Viets, 1926, Zool. Anz., 69: 196.

Genus *Calonyx* Walter

Generotype: *Partnunia lauta* Walter, 1906, Zool. Anz., 30: 571-572.

Calonyx Walter, 1907, op. cit., 31: 298.

Protzia Marshall, 1931, Trans. Wis. Acad. Sci., 26: 312-313.

Sporadoporus Wolcott, 1905, Trans. Amer. Micros. Soc., 26: 191.

constans Marshall, 1943, op. cit., 62: 323, pl. 4, figs. 30-34. [Marshall Collection]

ovata Marshall, 1931, Trans. Wis. Acad. Sci., 26: 312-313, pl. 7, figs. 9-12. [Marshall Collection]

Genus *Partnuniella* Viets

Generotype: *Partnuniella thermalis* Viets.

Partnuniella Viets, 1938, Verhandl. Int. Verein. Limnol., 8, (Tome 3), p. 214.

thermalis Viets, 1938, op. cit., pp. 215-219, figs. 1-5. [Marshall Collection]

var. *paucipora* Viets, 1938, op. cit., p. 219, figs. 6-7. [Marshall Collection]

Family CLATHROSPERCHONIDAE Lundblad

Clathrosparchonidae Lundblad, 1936, Zool. Anz., 115: 30.

Genus *Clathrosparchon* Lundblad

Generotype: *Clathrosparchon crassipalpis* Lundblad, 1936, op. cit., p. 30.

Clathrosparchon Lundblad, 1936, op. cit., p. 30.

americanus Habeeb, 1953, Leaflet Acad. Biol., no. 1, p. 1, figs. 1-5.

Family **HYDRYPHANTIDAE** Thor*Hydryphantidae* Thor, 1900, *Nyt. mag. naturvid.*, 38: 263.Genus **Hydryphantes** KochGenerotype: *Acarus ruber* Geer.*Hydryphantes* Koch, 1842, *Übers. Arachnidensystems*, 3: 30.**multioporus** Marshall, 1930, *Trans. Wis. Acad. Sci.*, 25: 247-248, pl. 5, figs. 1-3. [Marshall Collection]**ruber** Geer, 1776, *Mem. hist. ins.*, 7: 141-146, pl. 9, figs. 3-9.*novatus* Viets, 1949, *Abh. naturw. Ver. Bremen*, 32: 311.*novus* Viets, 1939, *Arch. Hydrobiol.*, 36: 75.var. *mozleyi* Marshall, 1929, *Univ. Toronto Stud.*, Biol. Ser., no. 33, pp. 68-69, pl. 2, figs. 12-13. [Marshall Collection]**tenuabilis** Marshall, 1926, *Univ. Iowa Stud. Nat. Hist.*, 11: 33, pl. 2, fig. 9, pl. 3, figs. 16-18. [Marshall Collection]Genus **Thyas** KochGenerotype: *Thyas venusta* Koch, 1836, *Deutschl. Crust., Myr., Arachn.*, Heft 5, no. 18.*Thyas* Koch, 1836, *op. cit.***stolli** Koenike, 1895, *Abh. naturw. Ver. Bremen*, 13: 194-196, pl. 2, figs. 29-32. [Marshall Collection]Genus **Marshallothyas** CookGenerotype: *Marshallothyas asopos* Cook, 1953, *Proc. Ent. Soc. Wash.*, 55: 306, figs. 1-10.*Marshallothyas* Cook, 1953, *op. cit.*, p. 305.**asopos** Cook, 1953, *op. cit.*, p. 306, figs. 1-10. [C.N.H.M. Collection]Genus **Panisus** KoenikeGenerotype: *Panisus michaeli* Koenike, 1896, *Zool. Anz.*, 19: 356-357.*Panisus* Koenike, 1896, *op. cit.*, p. 356.**cataphracta** Koenike, 1895, *Abh. naturw. Ver. Bremen*, 13: 196-198, pl. 2, figs. 33-35.Genus **Lundbladia** VietsGenerotype: *Lundbladia feuerborni* Viets, 1929, *Zool. Anz.*, 86: 50-52.*Lundbladia* Viets, 1929, *op. cit.*, pp. 49-50.**musciicola** Mitchell, 1953, *Amer. Midl. Nat.*, 49: 162-165, 15 figs. [C.N.H.M. Collection]Genus **Panisopsis** VietsGenerotype: *Thyas vigilans* Piersig, 1896, *Zool. Anz.*, 19: 441.*Panisopsis* Viets, 1926, *op. cit.*, 66: 147.**pedunculata** Koenike, 1895, *Abh. naturw. Ver. Bremen*, 13: 192-194, pl. 1, figs. 24-28. [Marshall Collection]

Family **THERMACARIDAE** Sokolow

Thermacaridae Sokolow, 1927, Zool. Anz., 73: 20.

Genus **Thermacarus** Sokolow

Generotype: *Thermacarus thermobius* Sokolow, op. cit., pp. 11-20, figs. 1-13.

Thermacarus Sokolow, 1927, op. cit., pp. 11-20.

nevadensis Marshall, 1928, Psyche, 35: 93-95, pl. 3, figs. 1-9. [Marshall Collection]

Family **HYDRODROMIDAE** Viets

Hydrodromidae Viets, 1936, Tierwelt Deutschlands, Teil 31-32, pp. 139-140.

Diplodontidae Lundblad, 1927, Zool. bidrag Uppsala, 11: 408.

Genus **Hydrodroma** Koch

Generotype: *Hydrachna despiciens* Müller.

Hydrodroma Koch, 1842, Übers. Arachnidensystems, 3: 32.

Diplodontus auct. (nec Dugés).

americanus Marshall, 1926, Univ. Iowa Stud. Nat. Hist., 11: 33, pl. 1, fig. 4, pl. 2, fig. 8, pl. 3, figs. 13-15. [Marshall Collection]

despiciens Müller, 1776, Zool. Danicae Prodromus, p. 190.

Family **PSEUDOHYDRYPHANTIDAE** Viets

Pseudohydrphantidae Viets, 1926, Zool. Anz., 69: 196.

Genus **Pseudohydrphant** Viets

Generotype: *Pseudohydrphant* *parvulus* Viets, 1907, Abh. naturw. Ver. Bremen, 19: 142-145, figs. 1-4.

Pseudohydrphant Viets, 1907, op. cit., p. 142.

latipalpus Marshall, 1924, Trans. Amer. Micros. Soc., 43: 245-246, pl. 14, figs. 44-45. [Marshall Collection]

orbicularis Marshall, 1929, Univ. Toronto Stud., Biol. Ser., no. 33, pp. 67-68, pl. 4, figs. 36-38. [Marshall Collection]

Family **TEUTONIIDAE** Lundblad

Teutoniidae Lundblad, 1927, Zool. bidrag Uppsala, 11: 409-410.

Genus **Teutonia** Koenike

Generotype: *Hygrobates cometes* Koch, 1837, Deutschl. Crust., Myr., Arachn., Heft 10, no. 23.

Teutonia Koenike, 1889, Zool. Anz., 12: 104.

lunata Marshall, 1924, Trans. Amer. Micros. Soc., 43: 245, pl. 14, figs. 48-50. [Marshall Collection]

Family SPERCHONIDAE Thor

Sperchonidae Thor, 1900, *Nyt. mag. naturvid.*, **38**: 264.

Genus *Sperchonopsis* Piersig

Generotype: *Sperchon verrucosus* Protz.

Sperchonopsis Piersig, 1897, *Sitzungs. naturf. Ges. Leipzig*, **22-23**: 52.

Pseudosperchon Marshall, 1933, *Trans. Amer. Micros. Soc.*, **52**: 37.

ovalis Marshall, 1929, *Univ. Toronto Stud., Biol. Ser.*, no. 33, pp. 66-67, pl. 4, figs. 31-33. [Marshall Collection]

verrucosa Protz, 1896, *Zool. Anz.*, **19**: 23-25, figs. 1-3.

Genus *Sperchon*¹ Kramer

Generotype: *Sperchon squamosus* Kramer, 1879, *Arch. Naturg.*, **45**, (Bd. 1), pp. 2-5, pl. 1, figs. 1a-1d.

Sperchon Kramer, 1877, *op. cit.*, **43**, (Bd. 1), p. 240.

crassipalpis Marshall, 1933, *Trans. Amer. Micros. Soc.*, **52**: 37, pl. 7, figs. 1-3. [Marshall Collection]

glandulosus Koenike, 1886, *Zeit. Wissen. zool.*, **43**: 279-284, pl. 9, figs. 17-24.

jasperensis Marshall, 1929, *Univ. Toronto Stud., Biol. Ser.*, no. 33, pp. 65-66, pl. 2, figs. 14-15. [Marshall Collection]

parmatum Koenike, 1895, *Abh. naturw. Ver. Bremen*, **13**: 202-204, pl. 2, figs. 40-41.

tenuipalpis Koenike, 1895, *op. cit.*, pp. 204-206, pl. 2, figs. 42-47.

Family LEBERTIIDAE Thor

Lebertiidae Thor, 1900, *Nyt. mag. naturvid.*, **38**: 264.

Genus *Lebertia* Neuman

Generotype: *Lebertia insignis* Neuman, 1880, *K. Svenska Vetenskap. Handl.*, n. ser., (4), **17**, (3), pp. 69-70, pl. 8, fig. 4.

Lebertia Neuman, 1880, *op. cit.*, pp. 68-69.

artaacetabula Marshall, 1912, *Trans. Amer. Micros. Soc.*, **31**: 227, pl. 27, figs. 1-3. [Marshall Collection]

distincta Marshall, 1914, *Trans. Wis. Acad. Sci.*, **17**: 1301, pl. 93, figs. 21-23. [Marshall Collection]

martisensis Marshall, 1943, *Trans. Amer. Micros. Soc.*, **62**: 408-410, pl. 3, figs. 23-25. [Marshall Collection]

needhami Marshall, 1943, *op. cit.*, p. 408, pl. 3, figs. 28-30. [Marshall Collection]

ontarioensis Marshall, 1929, *Univ. Toronto Stud., Biol. Ser.*, no. 33, pp. 70-71, pl. 5, figs. 46-47. [Marshall Collection]

¹ Four new forms of *Sperchon*, *S. brevirostris* ssp. *scabriosus*, *S. mitchelli*, *S. glandulosus canadensis*, and *S. plumifer acadiensis*, are proposed by Habeeb (*Leaf. Acad. Biol.*, no. 1, p. 8). Each form is described in less than fifty words and unillustrated. These descriptions do not satisfy the requirements of Article 25c of the International Code of Zoological Nomenclature.

- parmata** Marshall, 1912, Trans. Amer. Micros. Soc., 31: 226-227, pl. 27, figs. 4-5, pl. 28, figs. 6-7. [Marshall Collection]
- porosa** Thor, 1900, Nyt. mag. naturvid., 38: 273.
- quinquemaculosa** Marshall, 1929, Proc. Indiana Acad. Sci., 38: 315, 317, figs. 5-8. [Marshall Collection]
- setosa** Koenike, 1912, Trans. Roy. Canad. Inst., 9: 294-295, pl. 2, fig. 20.
tau-insignita Koenike, 1895, Abh. naturw. Ver. Bremen, 13: 201-202 (in part).
- tyrrelli** Koenike, 1912, Trans. Roy. Canad. Inst., 9: 290-293, pl. 1, fig. 16, pl. 2, figs. 13-15.
tau-insignita Koenike, 1895, Abh. naturw. Ver. Bremen, 13: 201-202 (in part).
- wolcottii** Koenike, 1912, Trans. Roy. Canad. Inst., 9: 292-294, pl. 1, figs. 17-18, pl. 2, fig. 19.
tau-insignita Koenike, 1895, Abh. naturw. Ver. Bremen, 13: 201-202 (in part).
- wyomingensis** Marshall, 1933, Trans. Amer. Micros. Soc., 52: 39-40, pl. 8, figs. 16-17. [Marshall Collection]

Genus *Oxus* Kramer

- Generotype: *Oxus oblongus* Kramer, 1879, Arch. Naturg., 45, (Bd. 1), pp. 5-7, pl. 1, figs. 2a-2c.
- Oxus* Kramer, 1877, op. cit., 43, (Bd. 1), pp. 240-241.
- connatus** Marshall, 1929, Univ. Toronto Stud., Biol. Ser., no. 33, pp. 71-72, pl. 4, fig. 35. [Marshall Collection]
- elongatus** Marshall, 1929, op. cit., p. 72, pl. 4, fig. 34.
- intermedius** Marshall, 1926, Univ. Iowa Stud. Nat. Hist., 11: 31, pl. 1, figs. 1-3. [Marshall Collection]

Genus *Frontipoda* Koenike

- Generotype: *Hydrachna musculus* Müller, 1776, Zool. Danicae Prodrömus, p. 191.
- Frontipoda* Koenike, 1891, Zool. Anz., 14: 19.
- americana** Marshall, 1914, Trans. Wis. Acad. Sci., 17: 1300-1301, pl. 92, figs. 1-8. [Marshall Collection]

Genus *Gnaphiscus* Koenike

- Generotype: *Gnaphiscus setosus* Koenike, 1898, Zool. Anz., 21: 262-263.
- Gnaphiscus* Koenike, 1898, loc. cit.
- occidentalis** Marshall, 1924, Trans. Amer. Micros. Soc., 43: 247, pl. 13, figs. 36-40, pl. 14, figs. 51-55. [Marshall Collection]

Family TORRENTICOLIDAE Oudemans

- Torrenticolidae* Oudemans, 1941, Zool. Anz., 136: 178.
- Atractidaeidae* Thor, 1902, op. cit., 25: 408.

Genus *Torrenticola* Piersig

Generotype: *Atractides anomalus* Koch, 1837, Deutschl. Crust., Myr., Arach., Heft 11, no. 10.

Torrenticola Piersig, 1897, Sitzungs. naturf. Ges. Leipzig, 22-23: 155.

Atractides auct. (nec Koch).

californicus Marshall, 1943, Trans. Amer. Micros. Soc., 62: 315-316, pl. 2, figs. 14-16. [Marshall Collection]

compactus Marshall, 1943, op. cit., p. 315, pl. 3, figs. 25-26, pl. 4, fig. 27. [Marshall Collection]

ellipsoidalis Marshall, 1943, op. cit., pp. 308-310, pl. 2, figs. 11-13. [Marshall Collection]

geographicus Marshall, 1943, op. cit., p. 314, pl. 3, figs. 21-24. [Marshall Collection]

indistinctus Marshall, 1929, Proc. Indiana Acad. Sci., 38: 317, 319, figs. 16-18. [Marshall Collection]

jordanensis Marshall, 1930, Trans. Wis. Acad. Sci., 25: 248-249, pl. 6, figs. 12-14. [Marshall Collection]

mercedensis Marshall, 1943, Trans. Amer. Micros. Soc., 62: 310, 312, pl. 2, figs. 9-10. [Marshall Collection]

oblongatus Marshall, 1943, op. cit., p. 316, pl. 4, figs. 28-29, pl. 5, fig. 40. [Marshall Collection]

obovatus Marshall, 1943, op. cit., pp. 316, 318, pl. 5, figs. 35-39. [Marshall Collection]

occidentalis Marshall, 1933, op. cit., 52: 40, pl. 7, figs. 5-7. [Marshall Collection]

sierrensis Marshall, 1943, op. cit., 62: 307, pl. 1, figs. 1-4. [Marshall Collection]

simulans Marshall, 1933, op. cit., 52: 40, pl. 7, figs. 8-10. [Marshall Collection]

tahoei Marshall, 1943, op. cit., 62: 308, pl. 1, figs. 5-7. [Marshall Collection]

waddelicus Marshall, 1943, op. cit., pp. 312, 314, pl. 3, figs. 17-20. [Marshall Collection]

Genus *Testudacarus* Walter

Generotype: *Testudacarus tripeltatus* Walter, 1928, Rec. Indian Mus., 30: pp. 75-78, figs. 8-10.

Testudacarus Walter, 1928, op. cit., p. 75.

americanus Marshall, 1943, Trans. Amer. Micros. Soc., 62: 320, pl. 6, figs. 46-48. [Marshall Collection]

minimus Marshall, 1943, op. cit., p. 322, pl. 6, figs. 43-45. [Marshall Collection]

Family LIMNESIIDAE Thor

Limnesiidae Thor, 1900, Nyt. mag. naturvid., 38: 265.

Genus *Limnesia* Koch

Generotype: *Limnesia fulgida* Koch.

Limnesia Koch, 1842, Übers. Arachnidensystems, 3: 27.

Limnesiopsis Piersig, 1897, Sitzungs. naturf. Ges. Leipzig, 22-23: 52.

- anomala** Koenike, 1895, Abh. naturw. Ver. Bremen, 13: 207-208, pl. 2, figs. 49-53.
subsp. *marshallae* Viets, 1938, Zool. Anz., 121: 134-135.
anomala Marshall, 1932, (male only), Trans. Wis. Acad. Sci., 27: 346, pl. 7, fig. 7.
- coerulea** Lundblad, 1941, Zool. Anz., 133: 158.
- columbica** Marshall, 1924, Trans. Amer. Micros. Soc., 43: 239-240, pl. 11, figs. 9-12. [Marshall Collection]
- cornuta** Wolcott, 1903, op. cit., 24: 89-92, pl. 12, figs. 1-5. [Marshall Collection]
- fulgida** Koch, 1835, Deutschl. Crust., Myr., Arachn., Heft 2, no. 19.
histrionica Wolcott, 1903, Trans. Amer. Micros. Soc., 24: 92-95, pl. 12, figs. 6-7.
wolcottii Piersig, 1905, Zool. Centr., 12: 198.
- koenikea** Piersig, 1894, Zool. Anz., 17: 151.
- maculata** Müller, 1776, Zool. Danicae Prodrum, p. 191.
americana Piersig, 1905, Zool. Centr., 12: 198.
elliptica Marshall, 1924, Trans. Amer. Micros. Soc., 43: 240, pl. 11, figs. 13-14. [Marshall Collection]
- marshallae** Viets, 1938, Zool. Anz., 121: 134-135.
anomala Marshall, 1927, Trans. Amer. Micros. Soc., 46: 213, pl. 7, figs. 5-6.
anomala Marshall, 1932, Trans. Wis. Acad. Sci., 27: 346, pl. 7, figs. 6-8.
hutchinsoni Lundblad, 1941, Zool. Anz., 133: 158.
- marshalliana** Lundblad, 1952, Ark. Zool., (2), 3: 525.
marshallae Lundblad, 1941, Zool. Anz., 133: 157.
- paucispina** Wolcott, 1903, Trans. Amer. Micros. Soc., 24: 98-100, pl. 13, figs. 10-11. [Marshall Collection]
- protractipora** Lundblad, 1941, Zool. Anz., 133: 157.
- undulata** Müller, 1781, Hydrachnae aquis Danicae, p. 80, pl. 11, fig. 1.
var. *californica* Lundblad, 1941, Zool. Anz., 133: 158.
- wawaseea** Marshall, 1929, Proc. Indiana Acad. Sci., 38: 315, figs. 9-11. [Marshall Collection]

Genus *Tyrrellia* Koenike

Generotype: *Tyrrellia circularis* Koenike.

- Tyrrellia* Koenike, 1895, Abh. naturw. Ver. Bremen, 13: 198-199.
circularis Koenike, 1895, op. cit., pp. 199-201, pl. 2, figs. 36-38.
var. *monensis* Marshall, 1940, Trans. Wis. Acad. Sci., 32: 386, pl. 1, figs. 5-7, pl. 2, fig. 15. [Marshall Collection]
- ovalis** Marshall, 1932, op. cit., 27: 342-343, pl. 8, figs. 18-21. [Marshall Collection]

Family PONTARACHNIDAE Thor

Pontarachnidae Thor, 1929, Nyt. mag. naturvid., 67, pl. 7.

Genus *Pontarachna* Philippi

Generotype: *Pontarachna punctulum* Philippi, 1840, Arch. Naturg., 6, (Bd. 1), pp. 191-193, pl. 4, figs. 4-5.

Pontarachna Philippi, 1840, op. cit., p. 193.

cruciata Hall, 1912, Pomona Coll., Rep. Laguna Mar. Lab., 1: 183-184, fig. 102.

Family HYGROBATIDAE Koch

Hygrobatides Koch, 1842, Übers. Arachnidensystems, 3: 7.

Genus *Hygrobates* Koch

Generotype: *Hygrobates lutescens* Koch, 1841, Deutschl. Crust., Myr., Arachn., Heft 37, no. 13.

Hygrobates Koch, 1837, op. cit., Heft 10, no. 8.

decaporus Koenike, 1895, Abh. naturw. Ver. Bremen, 13: 215-216, pl. 3, figs. 62-63.

exilis Koenike, 1895, op. cit., pp. 213-215, pl. 3, figs. 60-61.

longipalpis Hermann, 1804, Mém. Aptérolog., p. 55, pl. 3, fig. 1, pl. 9, fig. P.

ruber Marshall, 1926, Univ. Iowa Stud. Nat. Hist., 11: 32, pl. 1, figs. 5-7.

multiporus Koenike, 1895, Abh. naturw. Ver. Bremen, 13: 216-217, pl. 3, fig. 64. [Marshall Collection]

neooctoporus Marshall, 1926, Univ. Iowa Stud. Nat. Hist., 11: 34. [Marshall Collection]

octoporus Marshall, 1924, Trans. Amer. Micros. Soc., 43: 238-239, pl. 11, figs. 1-5.

occidentalis Marshall, 1943, op. cit., 62: 414, pl. 4, figs. 39-41. [Marshall Collection]

Genus *Atractides* Koch

Generotype: *Atractides spinipes* Koch, 1837, Deutschl. Crust., Myr., Arachn., Heft 11, no. 16.

Atractides Koch, 1837, loc. cit.

Megapus Neuman, 1880, K. Svenska Vetenskap. Handl., n. ser., (4), 17, (3), pp. 63-64.

crassipalpis Koenike, 1909, Abh. naturw. Ver. Bremen, 19: 246.

ovalis Koenike, 1895, op. cit., 13: 211-212, pl. 3, figs. 58-59.

nodipalpis Thor, 1899, Arch. Math. Naturv., 21, (5), pp. 39-40, pl. 17, figs. 120-121 (Palearctic).

var. *americanus* Marshall, 1943, Trans. Amer. Micros. Soc., 62: 414-415, pl. 4, figs. 31-34. [Marshall Collection]

spinipes Marshall, 1933, op. cit., 52: 41, pl. 7, figs. 18-20.

parviscutus Marshall, 1915, Trans. Amer. Micros. Soc., 34: 186, pl. 6, figs. 1-4. [Marshall Collection]

orthopes Marshall, 1915, op. cit., p. 187, pl. 6, figs. 10-12.

?*phenoplectes* Marshall, 1915, op. cit., pp. 186-187, pl. 6, figs. 5-9. [Marshall Collection]

sturgeonsis Marshall, 1927, op. cit., 46: 274-275, pl. 8, figs. 16-19. [Marshall Collection]

Family UNIONICOLIDAE Oudemans

Unionicolidae Oudemans, 1909, Tijdschr. Ent., 52: 60.

Genus *Unionicola* Haldeman

Generotype: *Unionicola crassipes* Müller.

Unionicola Haldeman, 1842, Zool. Contr., 1: 1.

Atax Koch, 1842, Übers. Arachnidensystems, 3: 7.

abnormipes Wolcott, 1898, Zool. Bull., 1: 280-281, fig. 2. [Marshall Collection]

aculeatus Koenike, 1890, Zool. Anz., 13: 139-140.

sayi Piersig, 1901, Das Tierreich, 13: 213.

adensameri Thon, 1901, Ann. Naturhist. Hofmus., 16: 31-35, pl. 3.

arcuata Wolcott, 1898, Zool. Bull., 1: 284-285, fig. 5. [Marshall Collection]

campelomaicola Marshall, 1935, Univ. Toronto Stud., Biol. Ser., no. 39, pp. 99-102, figs. 1-5. [Marshall Collection]

crassipes Müller, 1776, Zool. Danicae Prodrum, p. 189.

figuralis Koch, 1836, Deutschl. Crust., Myr., Arachn., Heft 7, no. 10.

fossulatus Koenike, 1895, Abh. naturw. Ver. Bremen, 13: 221-222, pl. 3, figs. 68-71.

indistinctus Wolcott, 1898, Zool. Bull., 1: 281-282, fig. 3. [Marshall Collection]

intermedia Koenike, 1882, Abh. naturw. Ver. Bremen, 7: 265-266 (Palearctic).

var. *wolcottii* Piersig, 1900, Zool. Anz., 23: 211-212. [Marshall Collection]

pectinatus Wolcott, 1898, Zool. Bull., 1: 280, fig. 1. [Marshall Collection]

serratus Wolcott, 1898, op. cit., pp. 282-283, fig. 4. [Marshall Collection]

stricta Wolcott, 1898, op. cit., pp. 283-284. [Marshall Collection]

tumidus Wolcott, 1898, op. cit., p. 285, fig. 6. [Marshall Collection]

ypsilophora Bonz, 1783, K. Deutsche Akad. Naturf. Halle, nova acta Leopoldana, 7: 52, pl. 1, figs. 1-4.

haldemani Piersig, 1900, Zool. Anz., 23: 212.

formosa Dana and Whelpley, 1836, Amer. Jour. Sci., (1), 30: 357-358, figs. 1-8.

Genus *Najadicola* Piersig

Generotype: *Unionicola ingens* Koenike.

Najadicola Piersig, 1897, Zool. Anz., 20: 59-60.

ingens Koenike, 1895, Abh. naturw. Ver. Bremen, 13: 219-220, pl. 3, figs. 65-67.

Genus *Neumania* Lebert

Generotype: *Hydrachna spinipes* Müller, 1776, Zool. Danicae Prodrum, p. 189.

Neumania Lebert, 1879, Bull. Soc. Vaud. Sci. Natur., 16: 357.

armata Marshall, 1922, Trans. Wis. Acad. Sci., 20: 209, pl. 3, figs. 11-13. [Marshall Collection]

muttkowskii Marshall, 1922, op. cit., p. 208, pl. 2, fig. 10 (in part). [Marshall Collection]

- distincta* Marshall, 1922, Trans. Wis. Acad. Sci., **20**: 210-211, pl. 4, figs. 20, 23. [Marshall Collection]
- brevibrachiata* Marshall, 1922, op. cit., p. 211, pl. 4, figs. 24-25. [Marshall Collection]
- okobojica* Marshall, 1926, Univ. Iowa Stud. Nat. Hist., **11**: 31, pl. 2, figs. 11-12. [Marshall Collection]
- extendens* Marshall, 1922, Trans. Wis. Acad. Sci., **20**: 209, pl. 3, figs. 14-16. [Marshall Collection]
- punctata* Marshall, 1922, (female only), op. cit., pp. 211-212, pl. 4, fig. 29. [Marshall Collection]
- fragilis* Marshall, 1922, op. cit., p. 208, pl. 2, figs. 7-9. [Marshall Collection]
- hickmani* Marshall, 1933, op. cit., **28**: 46, pl. 4, figs. 41-43. [Marshall Collection]
- kodiakica* Marshall, 1924, Trans. Amer. Micros. Soc., **43**: 246, pl. 14, figs. 42-43. [Marshall Collection]
- longiseta* Marshall, 1924, op. cit., p. 243, pl. 12, fig. 23, pl. 13, fig. 26. [Marshall Collection]
- ovata* Marshall, 1922, Trans. Wis. Acad. Sci., **20**: 206-207, pl. 2, fig. 1.
- papillator* Marshall, 1922, op. cit., pp. 209-210, pl. 3, figs. 17-20, pl. 4, fig. 21. [Marshall Collection]
- pubescens* Marshall, 1929, Proc. Indiana Acad. Sci., **38**: 319, figs. 1-4. [Marshall Collection]
- punctata* Marshall, 1922, (male only), Trans. Wis. Acad. Sci., **20**: 211-212. [Marshall Collection]
- semicircularis* Marshall, 1922, op. cit., p. 207, pl. 2, figs. 2-3. [Marshall Collection]
- tenuipalpis* Marshall, 1922, op. cit., pp. 207-208, pl. 2, figs. 4-6. [Marshall Collection]
- muttkowskii* Marshall, 1922, op. cit., p. 208, pl. 2, fig. 10 (in part). [Marshall Collection]
- ?*vernalis* Koenike, 1895, Abh. naturw. Ver. Bremen, **13**: 218-219.

Genus *Koenikea* Wolcott

Generotype: *Koenikea concava* Wolcott (male only).

Koenikea Wolcott, 1900, Trans. Amer. Micros. Soc., **21**: 189-190.

Tanaognathus Wolcott, 1900, op. cit., pp. 193-194.

- alata* Lundblad, 1943, K. Svenska Vetenskap. Handl., **20**, (5), p. 15, fig. 7.
- concava* Wolcott, 1900, (male only), Trans. Amer. Micros. Soc., **21**: 190-193, pl. 11, figs. 15, 17, 19, pl. 12, fig. 24. [Marshall Collection]
- haldemani* Viets, 1930, Zool. Anz., **92**: 268-272, figs. 6-8.
- marshallae* Viets, 1930, op. cit., pp. 267-271, figs. 1-5.
- spinipes* Wolcott, 1900, Trans. Amer. Micros. Soc., **21**: 194-197, pl. 12, figs. 25-28. [Marshall Collection]
- wolcottii* Viets, 1930, Zool. Anz., **92**: 266. [Marshall Collection]
- concava* Wolcott, 1900, (female only), Trans. Amer. Micros. Soc., **21**: 190-193, pl. 11, figs. 18, 20-21, pl. 12, fig. 23. [Marshall Collection]

Genus *Huitfeldtia* ThorGenerotype: *Huitfeldtia rectipes* Thor.*Huitfeldtia* Thor, 1898, Arch. Math. Naturv., **20**, (7), pp. 1-2.*rectipes* Thor, 1898, op. cit., pp. 4-5, pl. 5, figs. 1-7.Family **FELTRIIDAE** Thor*Feltriidae* Thor, 1929, Nyt. mag. naturvid., **67**, pl. 7.Genus *Feltria* KoenikeGenerotype: *Feltria minuta* Koenike.*Feltria* Koenike, 1892, Zool. Anz., **15**: 323.*minuta* Koenike, 1892, op. cit., pp. 323-324, figs. 3-4.Family **PIONIDAE** Thor*Pionidae* Thor, 1900, Nyt. mag. naturvid., **38**: 265.Genus *Wettina* PiersigGenerotype: *Tiphys podagricus* Koch, 1837, Deutschl. Crust., Myr., Arachn., Heft 11, no. 9.*Wettina* Piersig, 1892, Zool. Anz., **15**: 408-410.*primaria* Marshall, 1929, Univ. Toronto Stud., Biol. Ser., no. 33, p. 83, pl. 4, figs. 39-41. [Marshall Collection]Genus *Hydrochoreutes* KochGenerotype: *Spio unguolata* Koch.*Hydrochoreutes* Koch, 1837, Deutschl. Crust., Myr., Arach., Heft 11, no. 11.*ungulatus* Koch, 1835, op. cit., Heft 5, no. 17.Genus *Tiphys* KochGenerotype: *Tiphys sagulatus* Koch, 1837, Deutschl. Crust., Myr., Arachn., Heft 11, no. 7.*Tiphys* Koch, 1835, op. cit., Heft 5, no. 19.*Acercus* Koch, 1842, Übers. Arachnidensystems, **3**: 23.*Laminipes* Piersig, 1901, Zool. Anz., **24**: 219-220.*diversus* Marshall, 1929, Univ. Toronto Stud., Biol. Ser., no. 33, pp. 83-84, pl. 5, figs. 48-50. [Marshall Collection]*simulans* Marshall, 1924, Trans. Amer. Micros. Soc., **43**: 244, pl. 14, figs. 46-47. [Marshall Collection]*torris* Müller, 1776, Zool. Danicae Prodrum, p. 191.var. *americanus* Marshall, 1937, Trans. Wis. Acad. Sci., **30**: 237, pl. 8, figs. 58-61.

Genus *Pionopsis* Piersig

Generotype: *Hydrachne lutescens* Hermann, 1804, Mém. Aptérolog., p. 57, pl. 6, fig. 7.

Pionopsis Piersig, 1894, Zool. Anz., 17: 215-216.

latilamellis Marshall, 1924, Trans. Amer. Micros. Soc., 43: 244, pl. 13, figs. 32-34. [Marshall Collection]

Genus *Pionacercus* Piersig

Generotype: *Pionacercus leuckarti* Piersig, 1894, Zool. Anz., 17: 213-214, figs. 1-2.

Pionacercus Piersig, 1894, op. cit., p. 213.

novus Marshall, 1924, Trans. Amer. Micros. Soc., 43: 244-245, pl. 13, fig. 35, pl. 14, fig. 41. [Marshall Collection]

Genus *Piona* Koch

Generotype: *Nesaea rosea* Koch, 1837, Deutschl. Crust., Myr., Arachn., Heft 10, no. 7.

Piona Koch, 1842, Übers. Arachnidensystems, 3: 13.

Curripes Koenike, 1891, Zool. Anz., 14: 19-20.

americana Marshall, 1929, Trans. Wis. Acad. Sci., 24: 401-404, pl. 9, figs. 1-9. [Marshall Collection]

carnea Koch, 1836, Deutschl. Crust., Myr., Arachn., Heft 8, no. 24.

neocarnea Marshall, 1924, Trans. Amer. Micros. Soc., 43: 241, pl. 12, figs. 15-17. [Marshall Collection]

conglobata Koch, 1836, Deutschl. Crust., Myr., Arachn., Heft 8, no. 9 (Pale-arctic).

var. *wisconsinensis* Marshall, 1935, Trans. Wis. Acad. Sci., 29: 284, pl. 10, figs. 43-47. [Marshall Collection]

constrictus Wolcott, 1902, Trans. Amer. Micros. Soc., 23: 222-226, pl. 30, figs. 14-18. [Marshall Collection]

coronis Wolcott, 1902, op. cit., pp. 208-210, pl. 29, figs. 1-2. [Marshall Collection]

crassus Wolcott, 1902, op. cit., pp. 246-249, pl. 33, figs. 57-60. [Marshall Collection]

debilis Wolcott, 1902, op. cit., pp. 234-235, pl. 32, figs. 44-46. [Marshall Collection]

exilis Wolcott, 1902, op. cit., pp. 210-213, pl. 29, figs. 3-8. [Marshall Collection]

guatemalensis Stoll, 1887, Biol. Centr.-Amer., 13: 11-12, pl. 10, figs. 2-2b, pl. 11, figs. 1-1f.

inconstans Wolcott, 1902, Trans. Amer. Micros. Soc., 23: 241-243, pl. 32, fig. 47, pl. 33, figs. 48-50. [Marshall Collection]

insularis Marshall, 1924, op. cit., 43: 243, pl. 12, fig. 24, pl. 13, figs. 30, 31. [Marshall Collection]

interrupta Marshall, 1929, Univ. Toronto Stud., Biol. Ser., no. 33, pp. 81-82, pl. 6, figs. 59-60. [Marshall Collection]

linguaplax Crowell, 1953, Amer. Midl. Nat., 50: 426-431, figs. 1-12. [C.N.H.M. Collection]

- medius** Wolcott, 1902, Trans. Amer. Micros. Soc., 23: 229-231, pl. 31, figs. 30-31. [Marshall Collection]
- nodata** Müller, 1776, Zool. Danicae Prodrömus, p. 191 (Palearctic).
var. *latigenitalia* Marshall, 1924, Trans. Amer. Micros. Soc., 43: 242, pl. 12, fig. 22, pl. 13, figs. 27-29. [Marshall Collection]
? *fuscatus* Koenike, 1895, Abh. naturw. Ver. Bremen, 13: 209.
- obturbans** Piersig, 1896, Zool. Anz., 19: 439-440.
- pugilis** Wolcott, 1902, Trans. Amer. Micros. Soc., 23: 213-216, pl. 29, figs. 9-12. [Marshall Collection]
- reighardi** Wolcott, 1902, op. cit., pp. 235-239, pl. 32, figs. 35-38. [Marshall Collection]
- rotunda** Kramer, 1879, Arch. Naturg., 45, (Bd. 1), p. 12, pl. 1, fig. 6.
- rubrapes** Marshall, 1924, Trans. Amer. Micros. Soc., 43: 241-242, pl. 12, figs. 18-21. [Marshall Collection]
- setiger** Wolcott, 1902, op. cit., 23: 243-246, pl. 33, figs. 51-56. [Marshall Collection]
- socialis** Marshall, 1930, op. cit., 49: 343-344, pl. 39, figs. 5-10. [Marshall Collection]
- spinulosus** Wolcott, 1902, op. cit., pp. 226-229, pl. 31, figs. 24-29. [Marshall Collection]
- triangularis** Wolcott, 1902, op. cit., 23: 220-222, pl. 31, figs. 32-33. [Marshall Collection]
- turgidus** Wolcott, 1902, op. cit., pp. 216-220, pl. 30, figs. 19-23. [Marshall Collection]
- wolcottii** Marshall, 1937, Trans. Wis. Acad. Sci., 30: 235-236, pl. 7, figs. 49-52. [Marshall Collection]

Genus *Forelia* Haller

Generotype: *Arrhenurus variegator* Koch, 1837, Deutschl. Crust., Myr., Arachn., Heft 12, no. 24.

Forelia Haller, 1882, Mitt. naturf. Ges. Bern, 1881, Heft 2, pp. 58-59.

Tiphys Wolcott, 1905, Trans. Amer. Micros. Soc., 26: 214-215.

liliacea Müller, 1776, Zool. Danicae Prodrömus, p. 190.

ovalis Marshall, 1929, Univ. Toronto Stud., Biol. Ser., no. 33, pp. 78-79, pl. 5, figs. 42-45. [Marshall Collection]

Family AXONOPSIDAE Viets

Axonopsidae Viets, 1929, Zool. Anz., 80: 169.

Genus *Albia* Thon

Generotype: *Albia stationis* Thon.

Albia Thon, 1899, Zool. Anz., 22: 100-101.

caerulea Marshall, 1927, Trans. Amer. Micros. Soc., 46: 277, pl. 9, figs. 25-27. [Marshall Collection]

? **stationis** Thon, 1899, Zool. Anz., 22: 101-102.

Genus **Brachypoda** Lebert

Generotype: *Hydrachna versicolor* Müller, 1776, Zool. Danicae Prodrömus, p. 191.

Brachypoda Lebert, 1879, Bull. Soc. Vaud. Sci. Natur., (2), 16: 340, 374.

Brachypoda Habeeb, 1953, Leaflet. Acad. Biol., no. 1, p. 9 (in part).

setosicauda Habeeb, 1953, op. cit., p. 12, figs. 26-27.

ssp. *acuticauda* Habeeb, 1953, op. cit., p. 12, fig. 28.

Genus **Axonopsis** Piersig

Generotype: *Hydrachna complanata* Müller, 1776, Zool. Danicae Prodrömus, p. 191.

Axonopsis Piersig, 1893, Zool. Anz., 16: 310.

setoniensis Habeeb, 1953, Leaflet. Acad. Biol., no. 1, p. 9, figs. 23-25.

Genus **Ljania** Thor

Generotype: *Ljania bipapillata* Thor.

Ljania Thor, 1898, Arch. f. Math. Naturv., 20, (13), p. 3.

bipapillata Thor, 1898, op. cit., p. 3 (Palearctic).

ssp. *purpurea* Habeeb, 1953, Leaflet. Acad. Biol., no. 1, p. 9.

Genus **Neoaxonopsis** Lundblad

Generotype: *Neoaxonopsis odontogaster* Lundblad, 1938, Zool. Anz., 122: 38-39.

Neoaxonopsis Lundblad, 1938, op. cit., p. 38.

Brachypoda Habeeb, 1953, Leaflet. Acad. Biol., no. 1, p. 9 (in part).

pilositarsa Habeeb, 1953, Leaflet. Acad. Biol., no. 1, p. 9, figs. 20-23.

unguitarsa Habeeb, 1953, op. cit., pp. 9, 12, figs. 29-30.

Genus **Aturus** Kramer

Generotype: *Aturus scaber* Kramer, 1875, Arch. Naturg., 43, (Bd. 1), pp. 309-310, pl. 8, fig. 3.

Aturus Kramer, 1875, op. cit., p. 309.

acadiensis Habeeb, 1953, Leaflet. Acad. Biol., no. 1, p. 15.

canadensis Habeeb, 1953, Natural. Canad., 80: 274-275.

deceptor Habeeb, 1953, op. cit., pp. 275-276.

droueti Habeeb, 1953, Leaflet. Acad. Biol., no. 1, p. 15.

estellae Habeeb, 1953, op. cit., p. 2, figs. 6-11.

formosus Habeeb, 1953, op. cit., p. 4, figs. 12-14, 18.

ssp. *pallidus* Habeeb, op. cit., p. 4.

mirabilis Piersig, 1897, Sitzungs. naturf. Ges. Leipzig, 22-23: 157.

ssp. *scaber* Koenike, 1895, Abh. naturw. Ver. Bremen, 13: 186-191, pl. 1, fig. 23.

projector Habeeb, 1953, Natural. Canad., 80: 275.

Family MIDEIDAE Viets

Mideidae Viets, 1929, Zool. Anz., **80**: 169.

Genus *Midea* Bruzelius

Generotype: *Hydrachna orbiculata* Müller, 1776, Zool. Danicae Prodrumus, p. 190.

Midea Bruzelius, 1854, Biskrif. ofver Hydrachnider, p. 35.

determina Marshall, 1929, Univ. Toronto Stud., Biol. Ser., no. 33, pp. 84-85, pl. 7, figs. 62-66. [Marshall Collection]

expansa Marshall, 1940, Trans. Wis. Acad. Sci., **32**: 138-139, pl. 4, figs. 26-30. [Marshall Collection]

Family MIDEOPSIDAE Thor

Mideopsidae Thor, 1928, Zeits. Morph. Okol. Tiere, **11**: 109.

Genus *Mideopsis* Neuman

Generotype: *Hydrachna orbicularis* Müller.

Mideopsis Neuman, 1880, K. Svenska Vetenskap. Handl., n. ser., (4), **17**, (3), p. 67.

americanus Marshall, 1940, Trans. Wis. Acad. Sci., **32**: 140-141, pl. 1, figs. 1-4. [Marshall Collection]

fibrosa Lundblad, 1941, Zool. Anz., **133**: 159-160.

lamellipalpis Lundblad, 1941, op. cit., p. 159.

orbicularis Müller, 1776, Zool. Danicae Prodrumus, p. 190.

rosea Lundblad, 1941, Zool. Anz., **133**: 159.

Genus *Xystonotus* Wolcott

Generotype: *Xystonotus asper* Wolcott.

Xystonotus Wolcott, 1900, Trans. Amer. Micros. Soc., **21**: 185-186.

asper Wolcott, 1900, op. cit., pp. 186-189, pl. 10, figs. 8-14. [Marshall Collection]

reelfootensis Hoff, 1944, Jour. Tenn. Acad. Sci., **19**: 234-238, pl. 1, figs. 1-10. [Marshall Collection]

Family KRENDOWSKIJIDAE Lundblad

Krendowskijidae Lundblad, 1930, Zool. bidrag Uppsala, **13**: 56-58.

Genus *Krendowskia* Piersig

Generotype: *Krendowskia latissima* Piersig, 1895, Zool. Anz., **18**: 147-148.

Krendowskia Piersig, 1895, op. cit., pp. 147-148.

similis Viets, 1931, op. cit., **93**: 42-44, figs. 12-14.

Genus *Geayia* Thor

Generotype: *Geayia venezuelae* Thor, 1897, Bull. Mus. d'Hist. Nat., 3: 12-13, figs. 1-6.

Geayia Thor, 1897, op. cit., pp. 11-12.

Krendowskia Wolcott, 1900, Trans. Amer. Micros. Soc., 21: 178-181.

Krendowskia Wolcott, 1905, op. cit., 26: 193.

ovata Wolcott, 1900, Trans. Amer. Micros. Soc., 21: 181-185, pl. 9, figs. 1-7. [Marshall Collection]

Family ARRENURIDAE Thor

Arrenuridae Thor, 1900, Nyt. mag. naturvid., 38: 266.

Genus *Arrenurus* Dugés

Generotype: *Arrenurus viridis* Dugés, 1834, Ann. Sci. Nat. (Zool.), (2), 1: 154-156, pl. 10, figs. 18-21.

Arrenurus Dugés, 1834, op. cit., pp. 154-156.

Steganaspis Wolcott, 1901, Trans. Amer. Micros. Soc., 22: 105-106.

Subgenus *Truncaturus* Thor

Generotype: *Arrenurus paluster* Thor, 1900, Nyt. mag. naturvid., 38: 380-382, pl. 8, figs. 25-27.

Truncaturus Thor, 1900, op. cit., pp. 379-380.

kenki Marshall, 1944, Amer. Midl. Nat., 31: 631, 633, pl. 1, figs. 1-2, 6. [Marshall Collection]

Subgenus *Micruracarus* Viets

Generotype: *Arrhenurus forpicatus* Neuman, 1880, K. Svenska Vetenskap. Handl., n. ser., (4), 17, (3), p. 90, pl. 6, figs. 2.

Micruracarus Viets, 1911, Zool. Anz., 38: 504.

acutus Marshall, 1908, Trans. Amer. Micros. Soc., 28: 92, pl. 8, figs. 14-16. [Marshall Collection]

bicaudatus Marshall, 1908, op. cit., p. 91, pl. 7, figs. 8-10. [Marshall Collection]

couleensis Lavers, 1945, op. cit., 64: 233-234, pl. 1, figs. 1-4. [Marshall Collection]

crenellatus Marshall, 1908, op. cit., 28: 90-91, pl. 8, figs. 11-13. [Marshall Collection]

infundibularis Marshall, 1908, op. cit., pp. 93-94, pl. 8, fig. 20, pl. 9, figs. 21-22. [Marshall Collection]

laticaudatus Marshall, 1908, op. cit., pp. 95-96, pl. 9, figs. 23-25. [Marshall Collection]

lyriger Marshall, 1908, op. cit., pp. 94-95, pl. 9, fig. 26, pl. 10, figs. 27-28. [Marshall Collection]

montifer Marshall, 1908, op. cit., pp. 96-97, pl. 10, figs. 29-31. [Marshall Collection]

- muttkowskii** Marshall, 1940, Trans. Wis. Acad. Sci., **32**: 144, pl. 4, figs. 33-35. [Marshall Collection]
- pseudosetiger** Marshall, 1921, Trans. Amer. Micros. Soc., **40**: 170, pl. 9, figs. 4-6. [Marshall Collection]
- setiger** Marshall, 1910, op. cit., **29**: 97-98.
- rotundus** Marshall, 1908, op. cit., **28**: 89-90, pl. 7, figs. 1-4, pl. 9, fig. 128. [Marshall Collection]
- scutulatus** Marshall, 1908, op. cit., p. 93, pl. 8, figs. 17-19. [Marshall Collection]
- setiger** Koenike, 1895, Abh. naturw. Ver. Bremen, **13**: 178-182, pl. 1, figs. 11-13.
- ovalis** Marshall, 1908, Trans. Amer. Micros. Soc., **28**: 90, pl. 7, figs. 5-7. [Marshall Collection]

Subgenus **Megaluracarus** Viets

Generotype: *Hydrachna globator* Müller, 1776, Zool. Danicae Prodrumus, p. 188.

Megaluracarus Viets, 1911, Zool. Anz., **38**: 504.

apetiolatus Piersig, 1904, Zool. Centr., **11**: 210.

corniger Marshall, 1903, Trans. Wis. Acad. Sci., **14**: 155-156, pl. 15, fig. 7, *a-e*.

aphelocercus Lavers, 1945, Trans. Amer. Micros. Soc., **64**: 248, pl. 4, figs. 33-38. [Marshall Collection]

belonocercus Lavers, 1945, op. cit., pp. 238-239, pl. 1, figs. 7-12. [Marshall Collection]

birgei Marshall, 1903, Trans. Wis. Acad. Sci., **14**: 158-159, pl. 16, fig. 10, *a-f*, pl. 17, fig. 10, *e*. [Marshall Collection]

capillatus Marshall, 1908, Trans. Amer. Micros. Soc., **28**: 101-102, pl. 12, figs. 43-45.

cardiacus Marshall, 1903, Trans. Wis. Acad. Sci., **14**: 153-154, pl. 17, fig. 5, *d-i*, pl. 18, fig. 5, *a-c*. [Marshall Collection]

cornicularis Marshall, 1908, Trans. Amer. Micros. Soc., **28**: 112-113, pl. 14, figs. 68-70. [Marshall Collection]

elevatus Marshall, 1914, Trans. Wis. Acad. Sci., **17**: 1301-1302, pl. 1, figs. 12-15, pl. 2, fig. 16.

elongatus Marshall, 1924, op. cit., **21**: 214, pl. 7, figs. 6-7, pl. 8, figs. 14-15. [Marshall Collection]

expansus Marshall, 1908, Trans. Amer. Micros. Soc., **28**: 107-108, pl. 13, figs. 53-55. [Marshall Collection]

invaginatus Lavers, 1945, op. cit., **64**: 246, 248, pl. 3, figs. 27-32. [Marshall Collection]

kincaidi Lavers, 1945, op. cit., pp. 242, 244, pl. 3, figs. 22-26, pl. 6, fig. 63. [Marshall Collection]

krameri Koenike, 1895, Abh. naturw. Ver. Bremen, **13**: 182-185, pl. 1, figs. 16-20.

laversi Marshall, 1944, Amer. Midl. Nat., **31**: 633, 635, pl. 2, figs. 7-11, pl. 3, fig. 17. [Marshall Collection]

longicaudatus Marshall, 1908, Trans. Amer. Micros. Soc., **28**: 111-112, pl. 14, figs. 65-67. [Marshall Collection]

mamilianus Marshall, 1908, op. cit., pp. 98-99, pl. 10, figs. 32-34, pl. 11, fig. 35. [Marshall Collection]

- manubriator** Marshall, 1903, Trans. Wis. Acad. Sci., 14: 151-152, pl. 15, fig. 3, *b*, pl. 16, fig. 3, *a*, *c*, *d*, pl. 17, fig. 3, *e*, *f*. [Marshall Collection]
- marshalli** Piersig, 1904, Zool. Centr., 11: 210.
globator Marshall, 1903, Trans. Wis. Acad. Sci., 14: 148-150, pl. 14, fig. 1, *a-g*, pl. 15, fig. 1, *e*.
- megalurus** Marshall, 1903, op. cit., pp. 150-151, pl. 14, fig. 2, *b*, *c*, *e*, *f*, pl. 15, fig. 2, *a*, *d*. [Marshall Collection]
- var. *intermedius* Marshall, 1940, op. cit., 32: 147-148, pl. 5, figs. 41-44. [Marshall Collection]
- morrisoni** Marshall, 1904, op. cit., 14: 523-524, pl. 40, fig. 2, *a-d*.
- parallellatus** Marshall, 1903, op. cit., 14: 154-155, pl. 16, fig. 6, *a-c*, pl. 17, fig. 6, *d*, pl. 18, fig. 6, *e*. [Marshall Collection]
- prominulus** Marshall, 1908, Trans. Amer. Micros. Soc., 28: 108-109, pl. 13, figs. 56-60. [Marshall Collection]
- pseudocaudatus** Piersig, 1905, Zool. Centr., 12: 185.
caudatus Marshall, 1904, Trans. Wis. Acad. Sci., 14: 521-523, pl. 40, fig. 1, *a-c*.
- pseudoconicus** Piersig, 1904, Zool. Centr., 11: 210.
conicus Marshall, 1903, Trans. Wis. Acad. Sci., 14: 158, pl. 18, fig. 9, *a-d*.
- pseudocylindratus** Piersig, 1904, Zool. Centr., 11: 210.
cylindratus Marshall, 1903, Trans. Wis. Acad. Sci., 14: 156-157, pl. 17, fig. 8, *a-d*.
- rawsoni** Marshall, 1929, Univ. Toronto Stud., Biol. Ser., no. 33, p. 86, pl. 7, figs. 67-70. [Marshall Collection]
- rectangularis** Marshall, 1908, Trans. Amer. Micros. Soc., 28: 110, pl. 14, figs. 61-63. [Marshall Collection]
- rheophilous** Lavers, 1945, op. cit., 64: 239, pl. 2, figs. 13-17, pl. 4, fig. 43. [Marshall Collection]
- scutuliformis** Marshall, 1908, op. cit., 28: 100, pl. 11, figs. 39-42. [Marshall Collection]
- semicircularis** Piersig, 1904, Zool. Centr., 11: 210.
securiformis Marshall, 1903, Trans. Wis. Acad. Sci., 14: 152-153, pl. 18, fig. 4, *a-c*.
- simulans** Marshall, 1921, Trans. Amer. Micros. Soc., 40: 169, pl. 10, figs. 17-21. [Marshall Collection]
- solifer** Marshall, 1908, op. cit., 28: 99, pl. 11, figs. 36-38. [Marshall Collection]
- tahoei** Marshall, 1910, op. cit., 29: 100-101, pl. 1, fig. 7, pl. 2, figs. 11-14, pl. 3, fig. 3. [Marshall Collection]
- uniformis** Marshall, 1921, op. cit., 40: 169, pl. 9, figs. 1-3. [Marshall Collection]
- wardi** Marshall, 1940, Trans. Wis. Acad. Sci., 32: 148, pl. 6, figs. 52-54. [Marshall Collection]

Subgenus **Arrenurus** *s. str.*

- americanus** Marshall, 1908, Trans. Amer. Micros. Soc., 28: 126-128, pl. 21, figs. 112-117. [Marshall Collection]
- var. *mucronatus* Lavers, 1945, op. cit., 64: 255-256, pl. 5, fig. 50. [Marshall Collection]

- amplus** Marshall, 1908, Trans. Amer. Micros. Soc., 28: 122-123, pl. 20, figs. 102-105. [Marshall Collection]
- auricularis** Lavers, 1945, op. cit., 64: 258, 260, pl. 5, figs. 51-53, pl. 6, figs. 60-62. [Marshall Collection]
- auris** Lavers, 1945, op. cit., pp. 260-261, pl. 6, figs. 55-58. [Marshall Collection]
- bleptopetiolatus** Cook, 1954, op. cit., 73: 48, 50, pl. 1, figs. 3, 7. [C.N.H.M. Collection]
- cascadensis** Lavers, 1945, op. cit., 64: 253, pl. 5, figs. 44-49. [Marshall Collection]
- compactilis** Marshall, 1908, op. cit., 28: 120, pl. 18, figs. 93-95. [Marshall Collection]
- dentipetiolatus** Marshall, 1908, op. cit., p. 117, pl. 16, fig. 81, pl. 17, figs. 82-83. [Marshall Collection]
- drepanophorus** Cook, 1954, op. cit., 73: 51-52, pl. 1, fig. 8, pl. 2, fig. 22, pl. 3, fig. 23. [C.N.H.M. Collection]
- falcicornis** Marshall, 1908, op. cit., 28: 121-122, pl. 19, figs. 96-98. [Marshall Collection]
- fissicorniformis** Cook, 1954, op. cit., 73: 50, pl. 1, figs. 2, 4. [C.N.H.M. Collection]
- fissicornis** Marshall, 1908, op. cit., 28: 130, pl. 22, figs. 125-127. [Marshall Collection]
- flabellifer** Marshall, 1908, op. cit., pp. 125-126, pl. 22, figs. 122-124. [Marshall Collection]
- gennadus** Cook, 1954, op. cit., 73: 54, pl. 3, figs. 27-28. [C.N.H.M. Collection]
- hungerfordi** Cook, 1954, op. cit., p. 52, pl. 2, figs. 14-16. [C.N.H.M. Collection]
- interpositus** Koenike, 1895, Abh. naturw. Ver. Bremen, 13: 176-178, pl. 1, figs. 6-10.
angustocaudatus Marshall, 1908, Trans. Amer. Micros. Soc., 28: 116, pl. 16, figs. 77-79. [Marshall Collection]
- laticornis** Marshall, 1908, op. cit., p. 122, pl. 19, figs. 99-101. [Marshall Collection]
- lautus** Koenike, 1895, Abh. naturw. Ver. Bremen, 13: 172-176, pl. 1, figs. 1-5.
- magnicaudatus** Marshall, Trans. Amer. Micros. Soc., 28: 123-124, pl. 20, figs. 106-108. [Marshall Collection]
- major** Marshall, 1908, op. cit., pp. 128-129, pl. 21, figs. 118-120, pl. 22, fig. 121. [Marshall Collection]
- maryellenae** Cook, 1954, op. cit., 73: 57, pl. 4, figs. 40, 44. [C.N.H.M. Collection]
- neosuperior** Cook, 1954, op. cit., p. 56, pl. 4, figs. 37-38. [C.N.H.M. Collection]
- pinguisomus** Cook, 1954, op. cit., pp. 44, 46, pl. 1, figs. 9, 12. [C.N.H.M. Collection]
- pistillatus** Marshall, 1908, op. cit., 28: 119-120, pl. 18, figs. 90-92. [Marshall Collection]
- planus** Marshall, 1908, op. cit., pp. 115-116, pl. 16, figs. 75-76. [Marshall Collection]
subsp. *ventropetiolatus* Lavers, 1945, op. cit., 64: 250-251, pl. 4, figs. 39-42, pl. 6, fig. 59. [Marshall Collection]
- platy-rotundo-cuspidator** Münchberg, 1951, Arch. Hydrobiol., 45: 383-388, figs. 1-8.

- pleopetiolatus* Marshall, 1944, Amer. Midl. Nat., 31: 635, 637, pl. 3, figs. 12-16. [Marshall Collection]
- pollictus* Marshall, 1910, Trans. Amer. Micros. Soc., 29: 106-107, pl. 1, figs. 9-10, pl. 3, figs. 25-28. [Marshall Collection]
- pseudosuperior* Cook, 1954, op. cit., 73: 55-56, pl. 4, figs. 41-42. [C.N.H.M. Collection]
- reflexus* Marshall, 1908, op. cit., 28: 117-118, pl. 17, figs. 84-86. [Marshall Collection]
- serratus* Marshall, 1919, op. cit., 38: 276, pl. 29, figs. 1-7. [Marshall Collection]
- superior* Marshall, 1908, op. cit., 28: 124-125, pl. 20, figs. 109-111. [Marshall Collection]
- tacomaensis* Marshall, 1924, Trans. Wis. Acad. Sci., 21: 213-214, pl. 7, figs. 1-5. [Marshall Collection]
- tetratumuli* Münchberg, 1953, Zool. Jahrb. (Syst.), 82: 51-56, figs. 1-9.
- trifoliatus* Marshall, 1908, Trans. Amer. Micros. Soc., 28: 115, pl. 15, figs. 72-74. [Marshall Collection]
- wallensis* Cook, 1954, op. cit., 73: 57-58, pl. 4, figs. 36, 43. [C.N.H.M. Collection]
- wolcotti* Marshall, 1908, op. cit., 28: 118-119, pl. 17, fig. 87, pl. 18, figs. 88-89. [Marshall Collection]

Incertae sedis

The names listed below represent species that cannot be entered in the modern taxonomic system because of the inadequacy of the descriptions. The species are listed in alphabetical order.

- Arrhenurus acerformis* Marshall, 1924, Trans. Wis. Acad. Sci., 21: 215, pl. 8, figs. 8, 9, 13. [Marshall Collection]
- Arrhenurus hirsutus* Marshall, op. cit., p. 215, pl. 8, figs. 10-12. [Marshall Collection]
- [This and the previous species of *Arrhenurus* are based only on females and cannot be properly entered into the generic system.]
- Hydrachna coccinea* Haldeman, 1843, Proc. Acad. Nat. Sci. Philadelphia, 1: 196. [Probably Hydryphantidae.]
- Hydrachna nebulosa* Haldeman, op. cit., p. 196. [Not *Hydrachna*.]
- Hydrachna pyriformis* Dana and Whelpley, 1836, Amer. Jour. Sci., (1), 30: 358-359, pl. 1, fig. 9. [Probably *Unionicola*.]
- Hydrachna scabra* Haldeman, 1843, Proc. Acad. Nat. Sci. Philadelphia, 1: 184. [Hydryphantes?]
- Hydrachna triangularis* Say, 1821, Jour. Acad. Nat. Sci. Philadelphia, 2: 59. [Probably *Unionicola*.]
- Hydrachna tricolor* Packard, 1871, Amer. Jour. Sci., (3), 1: 108. [There is no clue to the genus of this species in the description.]
- Hydrachna 5-undulata* Haldeman, 1843, Proc. Acad. Nat. Sci. Philadelphia, 1: 184. [*Limnesia*?]
- Steganaspis arrenuroides* Wolcott, 1901, Trans. Amer. Micros. Soc., 22: 105-109, pl. 21, figs. 1-6. [A larval description that cannot be placed in a taxonomic system that is based on adult characters.]
- Unionicola humerosa* Haldeman, 1842, Zool. Contr., 1, (1), p. 2, pl. 1, fig. 11.
- Unionicola lactea* Haldeman, 1842, op. cit., p. 1, pl. 1, figs. 6-8.

- Unionicola lugubris* Haldeman, 1842, Zool. Contr., 1, (1), p. 2.
Unionicola oriformis Haldeman, 1842, op. cit., p. 1, pl. 1, figs. 1-5.
Unionicola personata Haldeman, 1842, op. cit., p. 2, pl. 1, fig. 10.
Unionicola proxima Haldeman, 1842, op. cit., p. 2.
Unionicola reticulata Haldeman, 1842, op. cit., p. 3, pl. 1, fig. 9.
Unionicola symmetrica Haldeman, 1842, op. cit., p. 2.
Unionicola unicolor Haldeman, 1842, op. cit., p. 3.

[The preceding nine species are all defined on color differences. I agree with Wolcott (1899, p. 233) that these species are likely to be *U. ypsilophora* Bonz.]

Rejected Records

Records of Old World species from the New World that cannot be verified or properly assigned are listed here.

- Leidy, 1883, Proc. Acad. Nat. Sci. Phila., 35: 44-46. *Atax* [*Unionicola*] *bonzi* Claparède is cited and probably represents an erroneous identification. There is no other record of the species from the New World.
 Crowell, 1952, Jour. Elisha Mitchell Sci. Soc., 68: 191-196, 2 pls. Seven Palearctic species are recorded for the first time from North America. Neither the illustrations nor the text gives significant specific features. I have examined the slides and cannot see specific characters.
Atractides (*Torrenticola*) *anomalus*, op. cit., p. 192.
Aturus intermedius, op. cit., p. 193.
Hygrobatas fluvialis, op. cit., pp. 192-193.
Lebertia natans, op. cit., p. 192.
Megapus (*Atractides*) *walleri*, op. cit., p. 193.
Sperchon longirostris, op. cit., p. 192.
Sperchon squamosus, op. cit., p. 191.
 Bergstrom, 1953, Trans. Amer. Micros. Soc., 72: 157-162. Forty species are listed, four of them recorded from North America for the first time. Many of the specimens on which these records are based have been examined and the technique of slide preparation is such that reliable specific identifications are impossible. I reject all the records but list only the species new to North America.
Eylais wilsoni, op. cit., p. 158.
Piona nodata, op. cit., p. 161.
Piona nodata laminata, op. cit., p. 161.
Piona uncata, op. cit., p. 161.

Additional Type Material in the Marshall Collection

The species listed below were described by Marshall from outside of North America and are represented by type material in the Marshall Collection.

- Arrenurus asiaticus* Marshall, 1919, Trans. Amer. Micros. Soc., 38: 276-277, pl. 29, figs. 8-10—China.
Arrenurus distinctus Marshall, 1919, op. cit., p. 277, pl. 30, figs. 14-16—China.
Arrenurus epimerosus Marshall, 1919, op. cit., pp. 278-279, pl. 31, figs. 25-28—Brazil.

- Arrenurus geei* Marshall, 1921, Trans. Amer. Micros. Soc., **40**: 172, pl. 11, figs. 26-29—China.
- Arrenurus habanicus* Marshall, 1927, Trans. Ill. Acad. Sci., **19**: 198—Cuba.
- Arrenurus maderius* Marshall, 1919, Trans. Amer. Micros. Soc., **38**: 279, pl. 31, figs. 29-32—Brazil.
- Arrenurus merrilli* Marshall, 1919, op. cit., pp. 277-278, pl. 30, figs. 17-18—Brazil.
- Arrenurus pisciscaudapetiolatus* Marshall, 1928, Trans. Wis. Acad. Sci., **23**: 605, pl. 15, figs. 20-25—China.
- Arrenurus quadricornicus* Marshall, 1919, Trans. Amer. Micros. Soc., **38**: 279-280, pl. 31, figs. 33-37—British Guiana.
- Arrenurus scapulatus* Marshall, 1910, op. cit., **29**: 98-99, pl. 1, figs. 2-6—Belgian Congo.
- Arrenurus soochowensis* Marshall, 1921, op. cit., **40**: 171, pl. 11, figs. 22-25—China.
- Arrenurus triconicus* Marshall, 1919, op. cit., **38**: 278, pl. 30, figs. 19-24—British Guiana.
- Arrenurus valenciis* Marshall, 1919, op. cit., p. 277, pl. 29, figs. 11-13—Venezuela.
- Hydrachna nova* Marshall, 1928, Trans. Wis. Acad. Sci., **23**: 601-602, pl. 13, figs. 1-4—China.
- Hydrachna simulans* Marshall, 1928, op. cit., p. 602, pl. 14, figs. 16-18—China.
- Koenikea indistincta* Marshall, 1936, Carnegie Inst. Wash. Pub., no. 457, p. 134, figs. 1-5—Yucatan.
- Limnesia koenikei* var. *asiatica* Marshall, 1928, op. cit., p. 604, pl. 14, figs. 13-15—China.
- Neumania cenotea* Marshall, 1936, Carnegie Inst. Wash. Pub., no. 457, pp. 133-134, figs. 6-8—Yucatan.
- Neumania geei* Marshall, 1928, Trans. Wis. Acad. Sci., **23**: 603-604, pl. 14, figs. 10-12—China.
- Piona erratica* Marshall, 1940, Trans. Amer. Micros. Soc., **59**: 377-378, pl. 2, figs. 8-11—Argentina.
- Piona marianaensis* Marshall, 1927, Trans. Ill. Acad. Sci., **19**: 198—Cuba.
- Piona pearsei* Marshall, 1936, Carnegie Inst. Wash. Pub., no. 457, pp. 134-135, figs. 9-14—Yucatan.
- Xystonotus torrei* Marshall, 1927, Trans. Ill. Acad. Sci., **19**: 198-199—Cuba.

BIBLIOGRAPHY

BAKER, EDWARD W., and WHARTON, G. W.

1952. An introduction to Acarology. 8+465 pp., 377 figs., 1 col. pl. Macmillan and Co., New York.

BERGSTROM, DAVID W.

1953. Hydracarina from the Rocky Mountain region. Trans. Amer. Micros. Soc., **72**: 157-162.

BERLESE, ANTONIO

1888. Acari Austro-Americani. Bull. Soc. Ent. Ital., **20**: 171-222, 9 pls.

BONZ, CHRISTOPHORI

1783. Observatio Dr. J. Christophori Gottlieb Bonz. K. Deutsche Akad. naturf. zu Halle, nova acta Leopold., **7**: 52-53, 1 pl.

BRUZELIUS, RAGNAR MAGNUS

1854. Biskrifning öfver Hydrachnider, som förekomma inom Skåne. Akademisk Afhandling. Inaugural dissertation submitted to the faculty of Lund. 66 pp., 5 pls. (Paper not seen.)

COOK, DAVID R.

1953. *Marshallothyas*, a new genus belonging to the subfamily Thyasinae. Proc. Ent. Soc. Wash., **55**: 305-308, 10 figs.

1954. Preliminary list of the Arrenuri of Michigan. Part I. The subgenus *Arrenurus*. Trans. Amer. Micros. Soc., **73**: 39-58, 4 pls.

CROWELL, ROBERT M.

1952. New records of rheophilic water-mites from northwestern North Carolina. Jour. Elisha Mitchell Sci. Soc., **68**: 191-196, 2 pls.

1953. Morphology and developmental stages of *Piona linguaplax* n. sp. (Hydracarina: Pionidae), a water-mite from North Carolina. Amer. Midl. Nat., **50**: 426-432, 14 figs.

DANA, JAMES D., and WHELPLEY, JAMES

1836. On two American species of the genus *Hydrachna*. Amer. Jour. Sci., (1), **30**: 354-359, 1 pl.

DUGÉS, ALFREDO

1884. *Atax alzatei*. La Naturaleza, **6**: 344-347, 1 pl.

DUGÉS, ANTONIO

1834. Deuxième Mémoire sur l'ordre des Acariens (1). Ann. Sci. Nat. (Zool.), (2), **1**: 144-174, 2 pls.

FABRICIUS, JOHANN C.

1805. Systema antiliatorum, . . . 14+372 pp. Brunsvigae. (Paper not seen.)

FAUST, ERNEST CARROLL

1918. Additions to our knowledge of *Unionicola aculeata* (Koenike). Trans. Amer. Micros. Soc., **37**: 125-128, 1 pl.

GEER, CHARLES DE

1752-78. *Memoires pour servir à l'histoire des insectes*. 1-8, 238 pls. Stockholm.

HABEEB, HERBERT

1950. Three interesting water-mites. *Natural. Canad.*, 77: 112-117, 2 pls.

1953a. North American Hydrachnellae, Acari. I-V. *Leaf. Acad. Biol.*, no. 1, pp. 1-16, 30 figs.

1953b. North American Hydrachnellae, Acari. VI. *Natural. Canad.*, 80: 274-277.

HALDEMAN, SAMUEL STEHMEN

1842. On some American species of Hydrachnidae. *Zool. Contr.*, 1, (1), 6 pp., 1 pl. Philadelphia.

1843a. Descriptions of two species of Entomostraca and two Hydrachnae. *Proc. Acad. Nat. Sci. Philadelphia*, 1: 184.

1843b. Description of two species of *Hydrachna* and one of *Daphnia*. *Op. cit.*, p. 196.

HALL, HARRY V. M.

1912. Some marine and terrestrial Acarina of Laguna beach. *Pomona Coll. Rep. Laguna Mar. Lab.*, 1: 177-186, 9 figs.

HALLER, G.

1882. Die Arten und Gattungen der schweizer Hydrachnidenfauna. *Mitt. naturf. Ges. Bern*, nos. 1020-28, 1881, Heft 2, pp. 18-834, 4 pls.

HERMANN, JOHANN FREDERICH

1804. *Mémoire aptérologique*. 6+144 pp., 9 col. pls. Strasbourg.

HOFF, C. CLAYTON

1942. Locality records of some Hydracarina from Illinois. *Trans. Ill. Acad. Sci.*, 35: 205-206.

1944a. A preliminary study of the Hydracarina of Reelfoot Lake, Tennessee. *Jour. Tenn. Acad. Sci.*, 19: 45-69, 3 pls.

1944b. A new species and additional records of water mites from Reelfoot Lake, Tennessee. *Op. cit.*, pp. 234-239, 1 pl.

HUMES, ARTHUR G., and JAMNBACK, HUGO A.

1950. A water-mite parasitic in fresh-water clams. *Psyche*, 57: 77-87, 4 figs. 1 table.

Koch, CARL LUDWIG

1835-41. *Deutschlands Crustaceen, Myriopoden, und Arachniden*. Hefte 1-40, 40 col. pls. Regensburg. (Paper not seen.)

1837-50. *Übersicht des Arachnidensystems*. 1-5. Nürnberg.

KOENIKE, F.

1882. Über das Hydrachniden-Genus *Atax* Fabr. *Abh. naturw. Ver. Bremen*, 7: 265-268.

1884. Verzeichniss von im Harz gesammelten Hydrachniden. *Op. cit.*, 8: 31-37.

1886. Zwei neue Hydrachniden aus dem Isergebirge. *Zeitschr. wissen. zool.*, 43: 277-284.

1889. Kurze Mittheilung über ein neues Hydrachniden-Genus. *Zool. Anz.*, 12: 103-104.

1890a. Eine neues Hydrachniden-Genus (*Teutonia*). *Arch. Naturg.*, 56, (Bd. 1), pp. 75-80, 1 pl.

- 1890b. Ein neuer Bivalven-Parasit. Zool. Anz., 13: 139-140.
- 1891a. Nomenclatorische Correctur innerhalb der Hydrachniden-Familie. Op. cit., 14: 19-20.
- 1891b. Kurzer Bericht über nordamericanische Hydrachniden. Op. cit., pp. 256-258.
1892. Zwei neue Hydrachniden-Gattung aus dem Rhätikon. Op. cit., 15: 320-324, 4 figs.
- 1895a. Nordamericanische Hydrachniden. Abh. naturw. Ver. Bremen, 13: 167-226, 3 pls.
- 1895b. Liste des Hydrachnides, recueillies par le Docteur Théod. Barrios en Palestine, en Syrie et en Égypte. Rev. biol. nord. Fr., 7: 139-147, 1 pl.
1896. Zwei neue Hydrachniden-Gattungen nebst sechs unbekannten-Arten. Zool. Anz., 19: 356-362.
1897. Zur Systematik der Gattung *Eylais* Latr. Abh. naturw. Ver. Bremen, 14: 279-295, 6 figs.
1898. Über *Oxus* Kram., *Frontipoda* Koen. und eine neue verwandte Gattung. Zool. Anz., 21: 262-263.
1899. Hydrachniden-Fauna von Madagaskar und Nossi-Bé. Abh. Senckenb. naturf. Ges., 21: 297-435, 10 pls.
1909. Beitrag zur Kenntnis der Hydrachniden. Abh. naturw. Ver. Bremen, 19: 217-266, 45 figs.
1912. A revision of my "Nordamericanische Hydrachniden." Trans. Roy. Canad. Inst., 9: 281-296, 2 pls.

KRAMER, P.

1875. Beiträge zur Naturgeschichte der Hydrachniden. Arch. Naturg., 41, (Bd. 1), pp. 263-332, 2 pls.
1877. Grundzüge zur Systematik der Milben. Op. cit., 43, (Bd. 1), pp. 215-247.
1879. Neue Acariden. Op. cit., 45, (Bd. 1), pp. 1-18, 2 pls.

LATREILLE, CITOYEN

1796. Précis des caractères, génériques des insectes, disposés dans un ordre naturel. 13+208 pp. Paris.

LAVERS, CHARLES H.

1941. A new species of *Limnochares* from North America. Univ. Wash. Pub., Biology, 12: 3-6, 1 pl.
1945. The species of *Arrenurus* of the state of Washington. Trans. Amer. Micros. Soc., 64: 228-264, 6 pls.

LEACH, WILLIAM ELFORD

1815. A tabular view of the external characters of four classes of animals, which Linné arranged under Insecta . . . Trans. Linn. Soc. London, 11: 306-400.

LEBERT, HERMANN

1879. Hydrachnides du Léman. Bull. Soc. Vaud. Sci. Natur., 16: 327-377, 2 pls.

LEIDY, J.

- 1883a. On the reproduction and parasites of *Anodonta fluviatilis*. Proc. Acad. Nat. Sci. Philadelphia, 35: 44-46.
- 1883b. On the parasites of *Anodonta fluviatilis*. Ann. Mag. Nat. Hist., (5), 11: 391-392. (Reprinting of previous paper.)

LINNAEUS, CAROLUS

1758. Systema naturae . . . 10th ed., 824 pp. Holmiae.

LUNDBLAD, O.

1927. Die Hydracarinen Schwedens I. Zool. bidrag Uppsala, 11: 185-540, 254 figs., 16 pls.
1930. Südamerikanische Hydracarinen. Op. cit., 13: 1-86, 42 figs., 3 pls.
1934. Die nordamerikanische Arten der Gattung *Hydrachna*. Ark. Zool., 28A, (3), 44 pp., 22 figs., 3 pls.
1936. Neue Wassermilben aus Santa Catharina in Südbrasilien. Zool. Anz., 115: 29-51.
1938. Neue brasilianische Wassermilben. Op. cit., 122: 35-44.
- 1941a. Neue Wassermilben aus Amerika, Afrika, Asien und Australien. Op. cit., 133: 155-160.
- 1941b. Eine Übersicht des Hydrachnellensystems und der jetzt bekannten Verbreitung der Gattungen dieser Gruppe. Zool. bidrag Uppsala, 20: 359-379.
- 1941-44. Die Hydracarinenfauna Südbrasilien und Paraguays. K. Svenska vetenskap. Handl., (3), 19, (7), 183 pp., 82 figs., 5 pls.; 20, (2), 175 pp., 80 figs., 7 pls.; 20, (5), 147 pp., 75 figs., 14 pls.; 20, (8), 171 pp., 78 figs., 15 pls.; 20, (13), 182 pp., 58 figs., 10 pls.
1949. Hydrachnellae. l'Inst. Parcs Nat. Congo Belge, Parc National Albert, II Mission H. Damas, fasc. 18, 87 pp., 50 figs., 8 pls.
1952. Hydracarinen von den Ostafrikanischen Gebirgen. Ark. Zool., (2), 3: 391-525, 74 figs., 16 pls.

MARSHALL, RUTH

1903. Ten species of *Arrhenuri* belonging to the subgenus *Megalurus* Thon. Trans. Wis. Acad. Sci., 14: 145-172, 5 pls.
1904. A new *Arrhenurus* and notes on the collections made in 1903. Op. cit., pp. 520-526, 1 pl.
1908. The *Arrhenuri* of the United States. Trans. Amer. Micros. Soc., 28: 85-140, 16 pls.
1910. New studies of the *Arrhenuri*. Op. cit., 29: 97-110, 3 pls.
1912. Some American *Lebertia*. Op. cit., 31: 225-230, 2 pls.
1914. Some new American water mites. Trans. Wis. Acad. Sci., 17: 1300-1304, 2 pls.
1915. American species of the genus *Atractides*. Trans. Amer. Micros. Soc., 34: 185-188, 1 pl.
1919. New species of water mites of the genus *Arrhenurus*. Op. cit., 38: 275-281, 3 pls.
1921. New species and collections of *Arrhenuri*. Op. cit., 40: 168-176, 3 pls.
1922. New American water mites of the genus *Neumania*. Trans. Wis. Acad. Sci., 20: 205-213, 3 pls.
1923. A preliminary report on the vertical distribution of water mites in Green Lake, Wisconsin. Trans. Ill. Acad. Sci., 15: 200-201.
- 1924a. *Arrhenuri* from Washington and Alaska. Trans. Wis. Acad. Sci., 21: 213-218, 2 pls.
- 1924b. A collecting trip to Alaska and the Canadian Northwest. Trans. Ill. Acad. Sci., 16: 51-53, 3 pls.
- 1924c. Water mites from Alaska and the Canadian Northwest. Trans. Amer. Micros. Soc., 43: 236-255, 4 pls.
1926. Water mites of the Okoboji region. Univ. Iowa Stud. Nat. Hist., 11: 28-35, 3 pls.
- 1927a. Collecting water mites in Cuba. Trans. Ill. Acad. Sci., 19: 197-199.
- 1927b. Water mites from Cuba. Trans. Amer. Micros. Soc., 46: 61-65, 2 pls.
- 1927c. Hydracarina of the Douglas Lake region. Op. cit., pp. 268-285, 3 pls.
- 1928a. Water mites from China. Trans. Wis. Acad. Sci., 23: 601-609, 3 pls.

- 1928b. A new species of water mite from thermal springs. *Psyche*, **35**: 92-97, 1 pl.
- 1929a. The water mites of Lake Wawasee. *Proc. Indiana Acad. Sci.*, **38**: 315-320, 2 pls.
- 1929b. The morphology and developmental stages of a new species of *Piona*. *Trans. Wis. Acad. Sci.*, **24**: 401-404, 1 pl.
- 1929c. Canadian Hydracarina. *Univ. Toronto Stud., Biol. Ser.*, no. 33 (Pub. Ont. Fish. Res. Lab., no. 39), pp. 57-93, 7 pls.
- 1930a. The water mites of the Jordan Lake region. *Trans. Wis. Acad. Sci.*, **25**: 245-253, 2 pls.
- 1930b. Hydracarina from Glacier National Park. *Trans. Amer. Micros. Soc.*, **49**: 342-345, 1 pl.
1931. Preliminary list of the Hydracarina of Wisconsin. Part I. The red mites. *Trans. Wis. Acad. Sci.*, **26**: 311-319, 2 pls.
1932. Preliminary list of the Hydracarina of Wisconsin. Part II. *Op. cit.*, **27**: 339-357, 4 pls.
- 1933a. Water mites from Wyoming as fish food. *Trans. Amer. Micros. Soc.*, **52**: 34-41, 2 pls.
- 1933b. Preliminary list of the Hydracarina of Wisconsin. Part III. *Trans. Wis. Acad. Sci.*, **28**: 37-61, 6 pls.
- 1935a. A new parasitic *Unionicola*. *Univ. Toronto Stud., Biol. Ser.*, no. 39 (Pub. Ont. Fish. Res. Lab., no. 53), pp. 97-101, 1 pl.
- 1935b. Preliminary list of the Hydracarina of Wisconsin. Part IV. *Trans. Wis. Acad. Sci.*, **29**: 273-297, 6 pls.
1936. Hydracarina from Yucatan. *Carnegie Inst. Wash. Pub.*, no. 457, pp. 133-137, 2 pls.
1937. Preliminary list of the Hydracarina of Wisconsin. Part V. *Trans. Wis. Acad. Sci.*, **30**: 225-251, 6 pls.
- 1940a. Water mites from Argentina. *Trans. Amer. Micros. Soc.*, **59**: 377-382, 2 pls.
- 1940b. On the occurrence of water mites in the food of turtles. *Amer. Midl. Nat.*, **24**: 361-364, 1 pl.
- 1940c. Preliminary list of the Hydracarina of Wisconsin. Part VI. *Trans. Wis. Acad. Sci.*, **32**: 135-165, 6 pls.
- 1940d. The water mite genus *Tyrrellia*. *Op. cit.*, pp. 383-389, 2 pls.
- 1943a. Hydracarina from California. Part I. *Trans. Amer. Micros. Soc.*, **62**: 306-324, 6 pls.
- 1943b. Hydracarina from California. Part II. *Op. cit.*, pp. 404-415, 4 pls.
1944. New species and notes on the Arrenuri. *Amer. Midl. Nat.*, **31**: 631-637, 3 pls.
1946. Preliminary list of the Hydracarina of Wisconsin. Revision of Part I. *Trans. Wis. Acad. Sci.*, **36**: 349-373, 5 pls.

MITCHELL, RODGER D.

1953. A new species of *Lundbladia* and remarks on the Hydryphantidae (water mites). *Amer. Midl. Nat.*, **49**: 159-170, 15 figs.
1954. A description of the water-mite, *Hydrovolzia gerhardi* new species, with observations on the life history and ecology. *Chicago Acad. Sci., Nat. Hist. Misc.*, no. 134, 9 pp., 3 figs.

MONTI, RINA

1905. Genere e specie nuovi de Idracnide. *Rend. Ist. Lomb. Sci. & Lett.*, (2), **38**: 163-176, 2 pls. (1 col.).

MOTAS, C.

1928. Contribution à la Connaissance des Hydracariens français particulièrement du Sud-Est de la France. Travaux Lab. Hydrobiol. Pisc., Univ. Grenoble, 20: 1-373, 247 figs., 9 photographs.

MÜLLER, OTTO FRIEDRICH

1776. Zoologiae Danicae Prodomus, seu animalium Daniae et Norvegiae indigenarum. 32+274 pp. Havniae.
1781. Hydrachnae, quas in aquis Daniae palustribus detexit, descripsit . . . 88 pp., 11 col. pls.

MÜNCHBERG, PAUL

1951. Ein zweiter Beitrag über die in den U. S. A. eine odanaten-parasitische Larvenphase aufweisenden *Arrenurus*-Arten (Ordnung: Hydracarina). Arch. Hydrobiol., 45: 378-388, 8 figs., 2 tables.
1952. Über Fortpflanzung, Lebensweise und Körperbau von *Arrenurus planus* Marsh. Zool. Jahrb., (Syst.), 81: 27-46, 12 figs.
1953. Vierter Beitrag zur Kenntnis der im nordamerikanischen Raume an Odonaten parasitierenden *Arrenurus*-Arten. Zool. Jahrb. (Syst.), 82: 47-57, 8 figs.

NEUMAN, CARL

1880. Om Sveriges Hydrachnider. K. Svenska Vetenskap. Handl., n. ser., (4), 17, (3), pp. 1-123, 14 pls.

OUDEMANS, A. C.

1909. Über die bis jetzt bekannten Thrombidium-Larven and über eine neue Klassifikation der Prostigmata. Tijdschr. Ent., 52: 19-60.
1926-37. Kritisch Historisch Overzicht der Acarologie. 1, 2, 3A-3G, illus. E. J. Brill, Leiden.
1941. Neue Funde auf dem Gebiete der Systematik und der Nomenklatur der Acari. VIII. Zool. Anz., 136: 177-186.

PACKARD, A. S., JR.

1871. On insects inhabiting salt water. Amer. Jour. Sci., (3), 1: 100-110, 5 figs.

PENNAK, ROBERT W.

1953. Fresh-water invertebrates of the United States. 9+769 pp., 470 figs. Hydracarina, pp. 470-487. New York.

PHILIPPI, A.

1840. Zoologische Bemerkungen. Arch. Naturg., 6, (Bd. 1), pp. 181-195, 2 pls.

PIERSIG, RICHARD

1892. Eine neue Hydrachniden-Gattung aus dem Sächsischen Erzgebirge. Zool. Anz., 15: 408-415, 3 figs.
1893. Neues über Wassermilben. Op. cit., 16: 309-312, 3 figs.
1894a. Über Hydrachniden. Op. cit., 17: 113-118, 7 figs.
1894b. Sachsens Wassermilben. Op. cit., pp. 213-216, 4 figs.
1895. Einiges über die Hydrachniden-Gattungen *Arrenurus* Dugés und *Thyas* C. L. Koch. Op. cit., 18: 145-148, 3 figs.
1896. Einige neue Hydrachniden-Formen. Op. cit., 19: 438-441.
1897a. Bemerkungen zur Hydrachnidenkunde. Op. cit., 20: 59-61.
1897b. Beiträge zur Kenntniss der in Sachsen einheimischen Hydrachniden-Formen. Sitzungs. naturf. Ges. Leipzig, 22-23, (1895-96), pp. 33-103.
1897c. Eine neue Hydrachniden-Gattung. Op. cit., pp. 155-157.

- 1897-1900. Deutschlands Hydrachniden. Zoologica, 22, Lief. 1, pp. 1-601. 51 pls.
1899. Einige neue *Eylais*-Arten. Zool. Anz., 22: 61-67, 8 figs.
1900. Hydrachnologische Bemerkungen. Op. cit., 23: 209-213.
1901a. Bemerkungen über die Gattung *Arrhenurus* Dugés. Op. cit., 24: 216-220.
1901b. Das Tierreich. Hydrachnidae und Halacaridae. 13, 18+336 pp., 87 figs.
1904. [Review of Marshall, 1903.] Zool. Centr., 11: 210-211.
1905a. [Review of Marshall, 1904.] Op. cit., 12: 185.
1905b. [Review of Wolcott, 1903.] Op. cit., pp. 197-198.

PROTZ, A.

1896. Beiträge zur Hydrachnidenkunde. Zool. Anz., 19: 23-26, 7 figs.

RADFORD, CHARLES D.

1950. Systematic check list of mite genera and type species. Union Internat. Sci. Biol., ser. C, no. 1, 232 pp.

SAY, THOMAS

1821. An account of the Arachnides of the United States. Jour. Acad. Nat. Sci. Philadelphia, 2: 59-81.

SOAR, CHARLES D., and WILLIAMSON, W.

- 1925-29. The British Hydracarina. Ray Society, 1, (1925), 10+216 pp., 20 pls.; 2, (1927), 7+215 pp., 20 pls.; 3, (1929), 8+184 pp., 20 pls.

SOKOLOW, IWAN

1927. *Thermacarus thermobius* n. gen. n. sp. eine Hydracarine aus heisser Quelle. Zool. Anz., 73: 11-20, 13 figs.

STOLL, OTTO

- 1886-93. Biologia Centrali-Americana, Zoology, Arachnida-Acarina. 21+55 pp., 21 pls.

SZALAY, L.

1949. Über die Hydracarinien der unterirdischen Gewässer. Hydrobiologia, 2: 141-179, 5 tables.

THON, KARL

1899. Ein neues Hydrachnidengenus aus Böhmen, nebst einigen Bemerkungen über böhmische *Hydryphantes*-Formen. Zool. Anz., 22: 100-102.
1901. Über eine neue parasitische *Atax*-Art aus Texas. Ann. Naturhist. Hofmus. (Vienna), 16: 31-35, 1 pl.

THOR, SIG

1897. Une intéressante Hydrachnidae nouvelle, provenant des récoltes de M. Geay au Venezuela. Bull. Mus. d'Hist. Nat., 3: 10-13, 6 figs.
1898. *Huitfeldtia* en ny Hydrachnide-slegt. Arch. Math. Naturv., 20, (7), pp. 1-6, 1 pl.
1899. Tredie bedrag til kundskaben om Norges Hydrachnider. Op. cit., 21, (5), pp. 1-64, 16 pls.
1900a. Prodrum systematicum Hydrachnidarum. Nyt. mag. naturvid., 38: 263-266.
1900b. Hydrachnologische notizen I-III. Op. cit., pp. 267-279, 2 pls.
1900c. Hydrachnologische notizen IV-VIII. Op. cit., pp. 369-388, 3 pls.
1902. Eigenartige bisher unbekannte Drüsen bei einzelnen "Hydrachniden" Formen. Zool. Anz., 25: 401-409, 5 figs.

1905. Eine interessante neue Milbengattung aus der schweizerischen Sammlung des Herrn Dr. W. Volz. Zool. Anz., 28: 505-509, 7 figs.
1916. Sur le genre *Hydrachna* Müll. et sur des nouvelles espèces provenant principalement de la Russie (Acarina, Hydrachnidae). Rev. Russe d'Ent., 16: 46-63, 20 figs.
1928. Untersuchung der von H. J. Stammer im Brachwasser des Parkteiches und des Ryckflusses bei Greifswald gesammelten Acarinen. Zeits. Morph. Okol. Tiere, 11: 104-114, 3 figs.
1929. Über die Phylogenie und Systematik der Acarina, mit Beiträgen zu ersten Entwicklungsgeschichte einzelner Gruppen. Nyt. mag. naturvid., 67: 145-210, 10 pls.

VIETS, KARL

1907. Neue Hydrachniden. Abh. naturw. Ver. Bremen, 19: 142-146, 3 figs.
1908. Eine neue *Thyas*-species. Zool. Anz., 33: 670-672, 2 figs.
1910. Hydrachnologische Beiträge III. Abh. naturw. Ver. Bremen, 20: 165-185, 16 figs.
1911. Ein Änderung in der Hydracarin-Nomenklatur. Zool. Anz., 38: 504.
- 1926a. Eine nomenclatorische Änderung im Hydrachniden-Genus *Thyas* C. L. Koch. Op. cit., 66: 145-148.
- 1926b. Versuch eines Systems der Hydracarin. Op. cit., 69: 188-199.
- 1929a. Einige neue Hydracarin-Gattungen aus Java und Bemerkungen zum System der Hydracarin. Op. cit., 80: 161-170.
- 1929b. Fünfte Mitteilung über Hydracarin von der Sunda-Inseln. Op. cit., 86: 49-56.
1930. Über nordamerikanische *Koenikea*-Arten (Hydracarina). Op. cit., 92: 266-272, 8 figs.
- 1931a. Über einige Gattungen und Arten der Axonopsae, Mideopsae, und Arrhenurae (Hydracarina). Op. cit., 93: 33-48, 14 figs.
- 1931b. Typus und Subgenera in *Hydrachna* O. F. Müller. Op. cit., pp. 173-185, 6 figs.
1933. Neue *Hydrachna*- und *Eylais*-Arten (Hydrachnellae, Acari) aus Porto Alegre, Brasilien. Op. cit., 103: 161-171, 11 figs.
1935. Die Wassermilben von Sumatra, Java, und Bali nach den Ergebnissen der Deutschen Limnologischen Sunda-Expedition. Ark. Hydrobiol., Suppl.-Bd. 13, pp. 494-798, 113 figs., Suppl.-Bd. 14, pp. 1-113, 18 figs., 3 pls., 1 map. [Also cited as parts 1-3 of Tropische Binnengewässer, 5.]
1936. Die Tierwelt Deutschlands und der angrenzenden Meersteile. Wassermilben oder Hydracarina. Teil 31-32, 7, 10+574 pp., 652 figs. Jena.
- 1938a. Über Hydrachnellae aus Uruguay. Zool. Anz., 121: 134-136, figs. 1-5.
- 1938b. Über die verschiedenen Biotope der Wassermilben, besonders über solche mit anormalen Lebensbedingungen und über einige neue Wassermilben aus Thermalgewässern. Verhandl. Int. Verein. theoret. angew. Limnol., 8, (Tome 3), pp. 209-224, 12 figs.
1939. Wassermilben (Hydrachnellae, Acari) aus den Bayrischen Alpen bei Garmisch-Partenkirchen. Arch. Hydrobiol., 36: 72-93, 19 figs.
1949. Nomenklatorische und taxonomische Bemerkungen zur Kenntnis der Wassermilben (Hydrachnellae, Acari). Abh. naturw. Ver. Bremen, 32: 292-327, 6 figs.
1953. Die aus Afrika bekannten Wassermilben (Hydrachnellae, Acari). Hydrobiologia, 5: 1-178.

VIETS, KURT O.

1949. Beitrag zur Kenntnis von *Eylais extendens* (O. F. Müller 1776) (Hydrachnellae, Acari). Veröffent. Mus. Natur-, Völker-, und Handelskunde in Bremen. Reihe A, Heft 1, pp. 41-105, illus.

VITZTHUM, H. GRAF

- 1940-43. Klassen und Ordnungen des Tierreichs. Acarina. 5, Abt. 4, Buch 5, Liefg. 1-7, 1011 pp., 522 figs. Leipzig.

WALTER, CHARLES

1906. Neue Hydrachnidenarten aus der Schweiz. Zool. Anz., 30: 570-575, 1 fig.
1907. Neue schweizerische Wassermilben. Op. cit., 31: 298-302, 1 fig.
1928. Zur Kenntnis der Mikrofauna von Britisch Indien. Hydracarina. Rec. Indian Mus., 30: 57-108, 43 figs.

WOLCOTT, ROBERT H.

1898. New American species of the genus *Atax* (Fab.) Bruz. Zool. Bull., 1: 279-285, 6 figs.
1899. On the North American species of the genus *Atax* (Fabr.) Bruz. Trans. Amer. Micros. Soc., 20: 193-259, 5 pls.
1900. New genera and species of North American Hydrachnidae. Op. cit., 21: 177-200, 4 pls.
1901. Description of a new genus of North American water mites, with observations on the classification of the group. Op. cit., 22: 105-117, 1 pl.
1902. The North American species of *Curvipes*. Op. cit., 23: 201-256, 5 pls.
1903. The North American species of *Limnesia*. Op. cit., 24: 85-107, 2 pls.
1905. A review of the genera of the water mites. Op. cit., 26: 161-243, 10 pls.
1918. The water-mites (Hydracarina). In Fresh-Water Biology, H. B. WARD and G. C. WHIPPLE, Editors. pp. 851-875, 34 figs. New York.

Index

Current names and principal references in roman type, synonyms in *italic* type

- abitibiensis Mars., 34
 abnormipes Wolc., 43
 acadiensis Hab. (Aturus), 48
 acadiensis Hab. (Limnoch.), 34
 acadiensis Hab. (Sperch.), 38
 Acarus L., 33, 34, 36
 Acercus Koch, 30, 45
 acerformis Mars., 54
 aculeatus Koen., 43
 acuticauda Hab., 48
 acutus Mars., 50
 adensameri Thon, 43
 alata Ldbl., 44
 Albia Thon, 47
 americana Ldbl. (Hydrach.), 34
 americana Ldbl. (Limnoch.), 34
 americana Mars. (Frontip.), 39
 americana Mars. (Hydrach.), 33
 americana Mars. (Piona), 46
 americana Piers. (Limnes.), 41
 americanus Hab. (Clath.), 35
 americanus Mars. (Arren.), 52
 americanus Mars. (Atract.), 42
 americanus Mars. (Hydro.), 37
 americanus Mars. (Mideop.), 49
 americanus Mars. (Testud.), 40
 americanus Mars. (Tiphys), 45
 amplus Mars., 53
 angustocaudatus Mars., 53
 anomala Koen. (Limnes.), 41
 anomala Mars. (Limnes.), 41
 anomalus Crow. (Torrent.), 55
 anomalus Koch (Torrent.), 40
 apetirolatus Piers., 51
 aphlocercus Lav., 51
 aquatica L., 34
 arcuata Wolc., 43
 armata Mars., 43
 ARRENURIDAE, 50
 arrenuroides Wolc., 54
 Arrenurus Dugés, 29, 30, 31, 47, 50, 52, 54, 55
 artaacetabula Mars., 38
 asiatica Mars., 56
 asiaticus Mars., 55
 asopos Cook, 36
 asper Wolc., 49
 Alax Koch, 43, 55
 ATTRACTIDAEIDAE, 30, 39
 Atractides auct., 30, 40, 55
 Atractides Koch, 30, 42, 55
 Aturus Kram., 48, 55
 auricularis Lav., 53
 auris Lav., 53
 AXONOPSIDAE, 47
 Axonopsis Piers., 48
 bakeri Mars., 34
 belonocercus Lav., 51
 bicaudatus Mars., 50
 bilunata Mars., 33
 bipapillata Thor, 48
 birgei Mars., 51
 bleptopetiolatus Cook, 53
 bonzi Leidy, 55
 Brachypoda Hab., 48
 Brachypoda Leb., 48
 brevibrachiata Mars., 44
 brevirostris Koen., 38
 caerulea Mars., 47
 californica Ldbl., 41
 californicus Mars., 40
 Calonyx Walt., 35
 campelomaicola Mars., 43
 canadensis Hab. (Aturus), 48
 canadensis Hab. (Sperch.), 38
 canadensis Mars. (Hydrach.), 34
 capillatus Mars., 51
 cardiacus Mars., 51
 carnea Koch, 46
 cascadiensis Lav., 53
 cataphracta Koen., 36
 caudatus Mars., 52
 cenotea Mars., 56
 Chitohydrachna Hab., 33
 circularis Koen., 41
 Clathrosperehon Ldbl., 35
 CLATHROSPERCHONIDAE, 35
 coccinea Hald., 54
 coerulea Ldbl., 41
 columbica Mars., 41
 cometes Koch, 37
 comosa Koen., 33
 compactilis Mars., 53
 compactus Mars., 40
 complanata Müll., 48
 concava Wolc., 44
 conglobata Koch, 46
 conicus Mars., 52
 conjecta Koen., 33

connatus Mars., 39
constans Mars., 35
constrictus Wolc., 46
cornicularis Mars., 51
corniger Mars., 51
cornuta Wolc., 41
coronis Wolc., 46
couleensis Lav., 50
crassipalpis Koen. (Atract.), 42
crassipalpis Ldbl. (Clath.), 35
crassipalpis Mars. (Sperch.), 38
crassipes Müll., 43
crassus Wolc., 46
crenellatus Mars., 50
crenulata Mars., 33
crinita Koen., 34
cruciata Hall, 42
cruenta Müll., 32, 33
Curvipes Koen., 46
cylindratus Mars., 52
Cyclothrix Wolc., 34

debilis Wolc., 46
decaporus Koen., 42
deceptor Hab., 48
dentipetiolatus Mars., 53
desecla Koen., 35
despiciens Dugés, 37
determina Mars., 49
diminuata Ldbl., 33
DIPLODONTIDAE, 37
Diplodontus auct., 37
Diplohydrachna Thor, 33
distincta Mars. (Leb.), 38
distincta Mars. (Neum.), 44
distinctus Mars. (Arren.), 55
diversus Mars., 45
dorsoscutata Viets, 33
drepanophorus Cook, 53
droueti Hab., 48

elevatus Mars., 51
ellipsoidalis Mars., 40
elliptica Mars., 41
elongatus Mars. (Arren.), 51
elongatus Mars. (Oxus), 39
ennishonensis Hab., 33
epimerosus Mars., 55
erratica Mars., 56
estellae Hab., 48
exilis Koen. (Hygro.), 42
exilis Wolc. (Piona), 46
expansa Mars., 49
expansus Mars., 51
extendens Müll. (Eyl.), 34
extendens Mars. (Neum.), 44
EYLAIDAE, 34
Eylais Latr., 34, 55

falcata Koen., 35
falcicornis Mars., 53

Feltria Koen., 45
FELTRIIDAE, 45
feuerborni Viets, 36
fibrosa Ldbl., 49
figuralis Koch, 43
fissicorniformis Cook, 53
fissicornis Mars., 53
5-undulata Hald., 54
flabellifer Mars., 53
fluvialis Crow., 55
forpicatus Neum., 50
Forelia Haller, 47
formosa D. and W., 43
formosus Hab., 48
fossulatus Koen., 43
fragilis Mars., 44
Frontipoda Koen., 39
fulgida Koch., 40, 41
fuscatus Koen. (Piona), 47

Geayia Thor, 50
geei Mars. (Arren.), 56
geei Mars. (Neum.), 56
gennadus Cook, 53
geographica Müll. (Hydrach.), 34
geographicus Mars. (Torrent.), 40
gerhardi Mitch., 32
gibberipons Viets, 35
glandulosus Koen., 38
globosa Geer, 33
globator Mars. (Arren.), 52
globator Müll. (Arren.), 51
Gnaphiscus Koen., 39
guatemalensis Stoll, 46

habanicus Mars., 56
haldemani Piers. (Union.), 43
haldemani Viets (Koen.), 44
hesperia Ldbl., 34
hickmani Mars., 44
hirsutipalpis Mars., 35
histrionica Wolc., 41
hirsutus Mars., 54
Huitfeldtia Thor, 45
humerosa Hald., 54
hungerfordi Cook (Arren.), 53
hungerfordi Ldbl. (Hydrach.), 33
hutchinsoni Ldbl. (Hydrach.), 33
hutchinsoni Ldbl. (Limnes.), 41
Hydrachna Leach, 29, 31, 32, 33, 34,
 37, 39, 43, 46, 48, 49, 51, 54, 56
HYDRACHNIDAE, 32
Hydrochoreutes Koch, 45
Hydrodroma Koch, 37
HYDRODROMIDAE, 37
Hydrovolzia Thor, 32
HYDROVOLZIIDAE, 32
Hydryphantes Koch, 36, 54
HYDRYPHANTIDAE, 36, 54
Hygrobatas Koch, 37, 42, 55
HYGROBATIDAE, 42

inconstans Wolc., 46
indistincta Mars. (Koen.), 56
indistinctus Mars. (Torrent.), 40
indistinctus Wolc. (Union.), 43
infundibularis Mars., 50
infundibulata Ldbl., 35
infundibulifera Koen., 35
ingens Koen., 43
insignis Neum., 38
insularis Mars., 46
intermedia Koen. (Union.), 43
intermedius Crow. (Aturus), 55
intermedius Mars. (Arren.), 52
intermedius Mars. (Oxus), 39
interpositus Koen., 53
interrupta Mars., 46
invaginatus Lav., 51

jasperensis Mars., 38
jordanensis Mars., 40

kenki Mars., 50
kincaidi Lav., 51
kodiakica Mars., 44
Koenikea Wolc., 44, 56
koenikea Wolc., 41
koenikei Piers. (Limnes.), 53
krameri Koen., 51
Krendowskia Piers., 49
Krendowskia Wolc., 50
 KRENDOWSKIJIDAE, 49

lactea Hald., 54
lamellipalpis Ldbl., 49
laminata Bergs., 55
Laminipes Piers., 45
laticaudatus Mars., 50
laticornis Mars., 53
latigenitalia Mars., 47
latilamellis Mars., 46
latipalpus Mars., 37
latissima Piers., 49
lauta Walt. (Calon.), 35
lautus Koen. (Arren.), 53
laversi Mars., 51
Lebertia Neum., 38, 55
 LEBERTIIDAE, 38
leuckarti Piers., 46
liliacea Müll., 47
Limnesia Koch, 40, 54, 56
 LIMNESIIDAE, 40
Limnestsopsis Piers., 40
Limnocharis Latr., 34
 LIMNOCHARIDAE, 34
linguaplex Crow., 46
Ljania Thor, 48
longicaudatus Mars., 51
longipalpis Herm. (Hygro.), 42
longirostris Crow., 55
longiseta Mars., 44
lugubris Hald., 55
lunata Mars., 37

Lundbladia Viets, 36
lutescens Koch (Hygro.), 42
lutescens Herm. (Pionops.), 46
lyriger Mars., 50

maculata Müll., 41
maderius Mars., 56
magnicaudatus Mars., 53
magniscutata Mars., 33
major Mars., 53
mamillanus Mars., 51
manubriator Mars., 52
marianaensis Mars., 56
marshallae Koen. (Eyl.), 35
marshallae Ldbl. (Hydrach.), 34
marshallae Ldbl. (Limnes.), 41
marshallae Viets (Koen.), 44
marshallae Viets (Limnes.), 41
marshalli Piers. (Arren.), 52
marshalliana Ldbl., 41
Marshallothyas Cook, 36
maryellenae Cook, 53
martisensis Mars., 38
medius Wolc., 47
Megaluracar Viets, 29, 51
megalurus Mars., 52
Megapus Neum., 30, 42, 55
mercedensis Mars., 40
merrilli Mars., 56
michaeli Koen., 36
microscutata Mars., 33
Micruracar Viets, 29, 50
Midea Bruz., 49
 MIDEIDAE, 49
 MIDEOPSIDAE, 49
Mideopsis Neum., 49
miliaria Berl., 33
minimus Mars., 40
minuta Koen., 45
mirabilis Piers., 48
mitchelli Hab., 38
monensis Mars., 41
montifer Mars., 50
morrisoni Mars., 52
mozleyi Mars., 36
mucronatus Lav., 52
multiaporus Mars. (Hydry.), 36
multiaporus Koen. (Hygro.), 42
musciola Mitch., 36
musculus Müll., 39
muttkowskii Mars. (Arren.), 51
muttkowskii Mars. (Neum.), 43, 44

Najadicola Piers., 43
natans Crow. (Leb.), 55
natans Lav. (Limnoch.), 34
nebulosa Hald., 54
needhami Mars., 38
Neoxonopsis Ldbl., 48
neocarnea Mars., 46
neooctoporus Mars., 42
neosuperior Mars., 53

- Nesaea* Koch, 46
Neumania Leb., 43, 56
nevadensis Mars., 37
nodata Bergs. (Piona), 55
nodata Müll. (Piona), 47
nodipalpis Thor, 42
nova Mars. (Hydrach.), 56
novatus Viets (Hydry.), 36
novus Viets (Hydry.), 36
novus Mars. (Pionac.), 46

oblongatus Mars., 40
oblongus Kram., 39
obovatus Mars., 40
obturbans Piers., 47
occidentalis Mars., 42
occidentalis Mars. (Gnaph.), 39
occidentalis Mars. (Torrent.), 40
octoporus Mars., 42
odontogaster Ldbl., 48
okobojica Mars., 44
ontarioensis Mars., 38
orbicularis Mars. (Pseudohy.), 37
orbicularis Müll. (Mideop.), 49
orbiculata Müll., 49
orthopes Mars., 42
ovalis Koen. (Atrac.), 42
ovalis Mars. (Arren.), 51
ovalis Mars. (Forel.), 47
ovalis Mars. (Sperch.), 38
ovalis Mars. (Tyrr.), 41
ovata Mars. (Calon.), 35
ovata Mars. (Neum.), 44
ovata Wolc. (Geayia), 50
oviformis Hald., 55
Oxus Kram., 39

pallidus Hab., 48
paluster Thor, 50
Panisopsis Viets, 36
Paniscus Koen., 36
papillator Mars., 44
parallellatus Mars., 52
parmata Mars. (Leb.), 39
parmatus Koen. (Sperch.), 38
Partnunia Piers., 35
Partnuniella Viets, 35
parviscutus Mars., 42
parvulus Viets, 37
paucipora Viets, 35
paucispina Wolc., 41
pearsei Mars., 56
pectinatus Wolc., 43
pedunculata Koen., 36
personata Hald., 55
phenoplectes Mars., 42
pilositarsa Hab., 48
pinguisomus Cook, 53
Piona Koch, 46, 55, 56
Pionacercus Piers., 46
PIONIDAE, 45
Pionopsis Piers., 46

pisciscaudapetiolatus Mars., 56
pistillatus Mars., 53
placophora Monti, 32
planus Mars., 53
platy-rotundo-cuspidator Münch., 53
pleopetiolatus Mars., 54
plumifer Thor, 38
podagricus Koch, 45
pollicitus Mars., 54
Polyxo Monti, 32
Pontarachna Phil., 42
PONTARACHNIDAE, 41
porosa Thor, 39
primaria Mars., 45
projector Hab., 48
prominulus Mars., 52
protractipora Ldbl., 41
Protzia Mars., 35
PROTZIIDAE, 35
proxima Hald., 55
pseudocaudatus Piers., 52
pseudoconicus Piers., 52
pseudocylindratus Piers., 52
Pseudohydraphantes Viets, 37
PSEUDOHYDRYPHANTIDAE, 37
pseudosetiger Mars., 51
Pseudosperchon Mars., 38
pseudosuperior Cook, 54
pubescens Mars., 44
pugilis Wolc., 47
punctata Mars., 44
punctulum Phil., 42
purpurea Hab., 48
pyriformis D. and W., 54

quadricornicus Mars., 56
quinquemaculosa Mars., 39

rawsoni Mars., 52
rectangularis Mars., 52
rectipes Thor, 45
reducta Ldbl., 33
reelfootensis Hoff, 49
reflexus Mars., 54
reighardi Wolc., 47
reticulata Hald., 55
Rhabdohydrachna Viets, 33
rheophilous Lav., 52
rimosa Piers., 35
robusta Mars., 35
rosea Ldbl. (Mideop.), 49
rosea Koch (Piona), 46
rotunda Mars. (Hydrac.), 33
rotunda Kram. (Piona), 47
rotundus Mars. (Arren.), 51
ruber Geer (Hydry.), 36
ruber Mars. (Hygro.), 42
rubrapes Mars., 47

sagulatus Koch, 45
sayi Piers., 43
scaber Kram. (Atur.), 48

- scaber* Koen. (Atur.), 48
scabra Hald., 54
scabriosus Hab., 38
scapulatus Mars., 56
Scutohydrachna Viets, 33
scutulatus Mars., 51
scutuliformis Mars., 52
securiformis Mars., 52
semicircularis Piers. (Arren.), 52
semicircularis Mars. (Neum.), 44
separata Ldbl., 33
serratus Mars. (Arren.), 54
serratus Wolc. (Union.), 43
setosicauda Hab., 48
setiger Koen. (Arren.), 51
setiger Mars. (Arren.), 51
setiger Wolc. (Piona), 47
setoniensis Hab., 48
setosa Koen. (Leb.), 39
setosus Koen. (Gnaph.), 39
sierrensis Mars., 40
similis Viets, 49
simulans Mars. (Arren.), 52
simulans Mars. (Hydrac.), 53
simulans Mars. (Tiph.), 45
simulans Mars. (Torrent.), 40
socialis Mars., 47
solifer Mars., 52
soochowensis Mars., 56
Sperchon Kram., 38, 55
SPERCHONIDAE, 38
Sperchonopsis Piers., 38
spinipes Koch (Atract.), 42
spinipes Mars. (Atract.), 42
spinipes Wolc. (Koen.), 44
spinipes Müll. (Neum.), 43
spinulosus Wolc., 47
Spio Koch, 45
Sporadoporus Wolc., 35
squamosus Crow. (Sperch.), 55
squamosus Kram. (Sperch.), 38
stationis Thon, 47
Steganaspis Wolc., 50, 54
stipata Ldbl., 34
stolli Koen., 36
stricta Wolc., 43
sturgeonensis Mars., 42
superior Mars., 54
symmetrica Hald., 55

tacomaensis Mars., 54
tahoei Mars. (Arren.), 52
tahoei Mars. (Torrent.), 40
Tanaognathus Wolc., 44
tau-insignita Koen., 39
tenuabilis Mars., 36
tenuipalpis Koen. (Sperch.), 38
tenuipalpis Mars. (Neum.), 44
Testudacarus Walt., 40
tetratumuli Münch., 54
Teutonia Koen., 37
TEUTONIIDAE, 37

Tetrahydrachna Ldbl., 33
Thyas Koch, 36
THERMACARIDAE, 37
Thermacarus Sok., 37
thermalis Viets, 35
thermobius Sok., 37
Tiphys Koch, 30, 45
Tiphys Wolc., 47
torrei Mars., 56
Torrenticola Piers., 30, 40, 55
TORRENTICOLIDAE, 30, 39
torris Müll., 45
triangularis Say (Hydrac.), 54
triangularis Wolc. (Piona), 47
triangulifera Koen., 35
tricolor Pack., 54
triconicus Mars., 56
trifolius Mars., 54
tripeltatus Walt., 40
Truncaturus Thor, 29, 50
tumidus Wolc., 43
turgidus Wolc., 47
tyrelli Koen., 39
Tyrrellia Koen., 41

uncata Bergs., 55
undulata Müll., 41
unguitarsa Hab., 48
ungulata Koch, 45
unicolor Hald., 55
uniformis Mars., 52
Unionicola Hald., 43, 54, 55
UNIONICOLIDAE, 43


valencius Mars., 56
variegator Koch, 47
venezuelae Thor, 50
ventropetiolatus Lav., 53
venusta Koch, 36
vernalis Koen., 44
verrucosa Protz., 38
versicolor Müll., 48
vigilans Piers., 36
viridis Dugés, 50

waddelicus Mars., 40
wallensis Cook, 54
walleri Crow., 55
wardi Mars., 52
wawassee Mars., 41
Wettina Piers., 45
wilsoni Bergs., 55
wisconsinensis Mars., 46
wolcottii Koen. (Leb.), 39
wolcottii Mars. (Arren.), 54
wolcottii Mars. (Piona), 47
wolcottii Piers. (Limnes.), 41
wolcottii Piers. (Union.), 43
wolcottii Viets (Koen.), 44
wyomingensis Mars., 39

Xystonotus Wolc., 49, 56
ypsilophora Bonz, 43, 55



UNIVERSITY OF ILLINOIS-URBANA



3 0112 027924213